

ReadiLink™ BSA Conjugation Kit

 Catalog number: 5501
 Unit size: 2 Reactions

Component	Storage	Amount (Cat No. 5501)
Component A: BSA (bovine serum albumin)	Refrigerated (2-8 °C), Minimize light exposure	2 X 2 mg
Component B: Conjugation Buffer (pH 4.7)	Refrigerated (2-8 °C), Minimize light exposure	1 bottle (20 mL)
Component C: EDC (1-ethyl 3-(3-dimethylaminopropyl) carbodiimide hydrochloride)	Refrigerated (2-8 °C), Minimize light exposure	2 X 10 mg
Component D: Purification Buffer Salts (pH 7.2)	Refrigerated (2-8 °C)	2 X 10 mL
Component E: Spin Desalting Columns (7K MWCO)	Refrigerated (2-8 °C)	2 X 2 mL

OVERVIEW

Bovine serum albumin (BSA) is a serum albumin protein that has numerous biochemical applications including ELISAs (Enzyme-Linked Immunosorbent Assay), immunoblots, and immunohistochemistry. Like most abundant plasma proteins, it is very stable and soluble. In addition, the 67 kDa protein is sufficiently large and complex to be fully immunogenic. It contains numerous sites per molecule for effective conjugation of peptides and other antigens using amine-reactive or carboxyl-reactive crosslinkers. Consequently, BSA is a popular carrier protein for conjugation to haptens and other weak antigens to make them more immunogenic for antibody production. This ReadiLink™ BSA Conjugation kit is primarily optimized for the simple preparation of hapten-carrier conjugates for immunization and antibody production. The ReadiLink™ BSA Conjugation kit is one-step conjugation of a hapten to a carrier protein using the carboxyl-reactive carbodiimide as the crosslinker. The resulting conjugate is used for eliciting an immune response and antibody production against the hapten. The carboxyl-reactive carbodiimide reacts with exposed carboxyl and amino groups on peptides and proteins to form stable bonds. These kits contain BSA formulated in buffers compatible with the carboxyl-reactive carbodiimide reactions and desalt spin columns, which offer exceptional protein recovery by simple centrifugation step.

AT A GLANCE
Protocol Summary

1. Prepare protein solution
2. Prepare hapten solution
3. Mix protein with hapten into EDC
4. Incubate the reaction at RT for 2 hrs
5. Purify the conjugate by desalting

Important

The following protocol is a general protocol for a wide variety of haptens. Optimize the protocol accordingly for the conjugation efficiencies based on the size and structure of your hapten. Using a molar excess of hapten over carrier protein ensures efficient conjugation. In general, a reaction with equal mass amounts of hapten and carrier protein will achieve sufficient molar excess.

SAMPLE EXPERIMENTAL PROTOCOL
Prepare BSA-Hapten Conjugation:

1. Add 200 µL of ddH₂O into the vial of BSA (Component A) to make

a 10 mg/mL solution.

2. Dissolve up to 2 mg hapten in 450 µL Conjugation Buffer (Component B).

Note: Some haptens might have limited solubility, use DMSO (<30% in the final conjugation solution) to dissolve it first. Higher concentrations of DMSO might irreversibly denature the carrier protein.

3. Add the 450 µL hapten solution into the 200 µL BSA solution to have Hapten-BSA mixture solution.
4. Add the Hapten-BSA solution into one vial of EDC (10mg) (Component C), dissolve it by gentle mixing. Incubate at room temperature for 2 hours.
5. Purify the conjugate by desalting to remove non-reacted crosslinker and protein preservative (e.g., sodium azide).

Purify BSA-Hapten conjugate:

1. Thaw 1 bottle of Purification Buffer (Component D) to room temperature before use. Unused buffer can be stored at 4 °C for 1 week.
2. Twist off the bottom closure of the desalting column (Component E) and loosen the cap. Place the column in a collection tube.
3. Centrifuge the column at 1,000g for 2 minutes to remove the storage solution.
4. Remove the cap and slowly add 1 mL of purification buffer to the column. Centrifuge at 1,000g for 2 minutes, remove the buffer. Repeat this step for 3 additional times, discarding the buffer from the collection tube.
5. Place the column to a new collection tube, and gently apply the sample into the center of the compact resin bed.
6. Centrifuge the column at 1,000g for 2 minutes to collect the sample.
7. The Hapten-BSA conjugate can now be used for immunization. If the Hapten-BSA conjugate will be stored for more than a few days, sterile filter and store at 4°C or -20°C.

Note: If the conjugate is to be used within one week, PBS may be used for purification. If the conjugate is to be frozen, use the purification buffer salts (Component D) for purification. If DMSO is used in the conjugation, prepare the purification buffer salts with the same percentage of DMSO used for conjugation. This will minimize the precipitation in the column during desalting. If a precipitate formed during conjugation, centrifuge the

precipitated material, collect the supernatant, and save the precipitate. Purify the supernatant. Combine the precipitate and the purified conjugate.

EXAMPLE DATA ANALYSIS AND FIGURES




Figure 1.

EDC reacts with a carboxyl group of carrier protein BSA or KLH (represented by the red ball), forming an amine-reactive O-acylisourea intermediate (the central molecule). The O-acylisourea intermediate reacts with an amine group on the antigen molecule represented by the smaller blue ball, yielding a conjugate of the two molecules joined by a stable amide bond [Please note the O-acylisourea intermediate is also susceptible to hydrolysis, making it unstable and short-lived in aqueous solution].

DISCLAIMER

AAT Bioquest provides high-quality reagents and materials for research use only. For proper handling of potentially hazardous chemicals, please consult the Safety Data Sheet (SDS) provided for the product. Chemical analysis and/or reverse engineering of any kit or its components is strictly prohibited without written permission from AAT Bioquest. Please call 408-733-1055 or email info@aatbio.com if you have any questions.