

(Z-LLE)2R110 Catalog number: 13466

Unit size: 1 mg

Product Details

Storage Conditions Freeze (<-15 °C), Minimize light exposure

Expiration Date 12 months upon receiving

Chemical Properties

Appearance Light yellow solid

Molecular Weight 1309.46

Soluble In DMSO

Chemical Structure

Spectral Properties

Excitation Wavelength 500 nm

Emission Wavelength 522 nm

Applications

The non-fluorescent R110 substrates generate the bright green fluorescent rhodamine 110 product that has Ex/Em = 494/521 nm, and can be easily detected with a FITC filter set. In general, R110 substrates are much more sensitive than the AMC-, AFC- or 4-nitroaniline-based substrates. This R110 substrate is used for monitoring the protease activities of the proteasome. The most common form of the proteasome is known as the 26S proteasome that contains one 20S core particle structure and two 19S regulatory caps. All 20S particles consist of four stacked heptameric ring structures that are themselves composed of two different types of subunits; alpha subunits are structural in nature, whereas beta subunits are predominantly catalytic. The outer two rings in the stack consist of seven alpha subunits each, which serve as docking domains for the regulatory particles and the alpha subunits N-termini form a gate that blocks unregulated access of substrates to the interior cavity. The inner two rings each consist of seven beta subunits and contain the protease active sites that perform the proteolysis reactions. AAT Bioquest offers a group of R110 substrates for monitoring the protease activities of the proteasome at different subsites, i.e., (i) sub-sites: beta1c, Z-LLE-R110; beta2c, Ac-KQL-R110; beta5c, Ac-WLA-R110; beta1i, Ac-PAL-R110; beta2i, Ac-KQL-R110; beta5c, Ac-WLA-R110 and Suc-LLVY-R110; and beta5i, Ac-ANW-R110. The protease activity is measured by monitoring the R110 liberation over time using excitation and emission wavelengths of 490 nm and 520 nm respectively.