

# XFD700 Anti-human CD11b Antibody \*HI11b, XFD700 Same Structure to Alexa Fluor™ 700\*

## **Product Details**

Storage Conditions 2-8°C with minimized light exposure. Do not freeze.

Expiration Date 12 months upon receiving

Concentration 0.1 mg/mL

Formulation Phosphate-buffered saline (PBS, pH 7.2), 0.09% sodium azide, 0.2% (w/v) BSA

#### **Antibody Properties**

Species Reactivity Human

Class Primary

Clonality Monoclonal

Host Mouse

Isotype Mouse IgG2b

Immunogen CD11b (CR3, Mac-1, Mo1, ITGAM, Integrin alpha-M)

Clone HI11b

Conjugate AF700

## **Biological Properties**

Appearance Dark blue liquid

Preparation Antibody purified by affinity chromatography and then conjugated with AF700 under optimal

conditions

Application Flow Cytometry (FACS), Fluorescence Imaging

### **Spectral Properties**

Conjugate AF700

Excitation Wavelength 696 nm

Emission Wavelength 719 nm

#### **Applications**

HI11b is an anti-human monoclonal antibody that recognizes the CD11b antigen. CD11b (sometimes referred to as Integrin  $\alpha$ M chain, C3biR, CR3A or CR3) is a 165 - 170 kD glycoprotein that is expressed on the surface of cells such as B cells, granulocytes, NK cells, macrophages and

dendritic cells. CD11b is involved with important cellular pathways, in particular, the cytokine-mediated signaling pathway, toll-like receptor 4 signaling pathway and apoptotic signaling pathway. Additionally, in many organisms, it is a promoter of superoxide anion generation, positively regulates microglial cell mediated cytotoxicity and is a positive regulator of prostaglandin-E synthase activity. CD11b has been associated with critical biological processes like cell adhesion, specifically cell-cell adhesion via plasma-membrane adhesion molecules, and is associated with a variety of biologically interesting macromolecules/ligands, for example, Factor X, iC3b and ICAM-1. CD11b is a very popular antibody target, with over 45000 publications in the last decade. CD11b has been widely used in innate immunity, neuroscience cell markers and cell biology research, often serving as a phenotypic marker for differentiating cell types in flow cytometric applications. This antibody was purified through affinity chromatography and conjugated to XFD700 (ex/em = 696/719 nm). XFD700 is manufactured by AAT Bioquest, and it has the same chemical structure of Alexa Fluor® 700 (Alexa Fluor® is the trademark of ThermoFisher).