

APC Anti-human CD138 Antibody *MI15*Catalog number: 113801C0, 113801C1, 113801C2
Unit size: 25 tests, 100 tests, 500 tests**Product Details**

Storage Conditions	2-8°C with minimized light exposure. Do not freeze.
Expiration Date	12 months upon receiving
Concentration	0.1 mg/mL
Formulation	Phosphate-buffered saline (PBS, pH 7.2), 0.09% sodium azide, 0.2% (w/v) BSA

Antibody Properties

Species Reactivity	Human
Class	Primary
Clonality	Monoclonal
Host	Mouse
Isotype	Mouse IgG1
Immunogen	CD138 (Syndecan-1)
Clone	MI15
Conjugate	APC

Biological Properties

Preparation	Antibody purified by affinity chromatography and then conjugated with APC under optimal conditions
Application	Flow Cytometry (FACS)

Spectral Properties

Conjugate	APC
Excitation Wavelength	651 nm
Emission Wavelength	660 nm

Applications

The MI15 monoclonal antibody reacts with human CD138, a 100 - 200 kD transmembrane protein typically expressed on the surface of b cells, plasma cells, epithelial cells and endothelial cells. CD138 is involved with important cellular pathways, in particular, the cytokine-mediated signaling pathway and canonical Wnt signaling pathway. Also, in certain organisms, it enhances extracellular exosome assembly and is a promoter of exosomal secretion. From a research standpoint, it is of biological interest due to its association with critical macromolecules/ligands such as Fibronectin. CD138 is a fairly uncommon antibody target, with a little more than 7000 publications in the last decade. Even still, CD138 is frequently used in flow cytometry applications as a phenotypic marker for differentiation of cell types, specifically in the study of cell motility/cytoskeleton/structure, cell biology and synaptic biology. This antibody was purified through affinity chromatography

and conjugated to APC (ex/em = 651/660 nm). It is compatible with the 642 nm laser and 702/85 nm bandpass filter (for example, as in the Luminex Amnis ImageStream).