

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no AS18 4211-1ml

Anti-Arabinogalactan-protein, AGP (monoclonal, clone LM2)

Product information

Immunogen Polysaccharide Arabinogalactan-protein (AGP) from *Oryza sativa*

Host Rat

Clonality Monoclonal

Subclass/isotype IgM

Purity Cell culture supernatant.

Format Liquid

Quantity 1 ml

Storage Store at +4°C (short term) and at -20°C (long term).

Additional information

Contains 0.05% Sodium Azide.

This antibody is made to rice arabinogalactan-proteins (AGPs) and it recognizes a carbohydrate epitope containing B-linked glucuronic acid.

In competitive inhibition ELISAs antibody binding to gum arabic was inhibited (50%) by 70 mg/ml 1-O-methyl-B-D-GlcA. The binding of the antibody to AGPs can be fully inhibited by 10 mM 1-O-methyl-B-D-GlcA.

Application information

Recommended dilution 1:10 (ELISA, IF)

Confirmed reactivity Higher plants, ferns and mosses

Not reactive in No confirmed exceptions from predicted reactivity are currently known

Additional information Antibody is recognizing carbohydrate epitope containing -linked glucuronic acid.

Selected references

Stacey et al. (1990). Patterns of expression of the JIM4 arabinogalactan-protein epitope in

Stacey et al. (1990). Patterns of expression of the JIM4 arabinogalactan-protein epitope in cell cultures and during somatic embryogenesis in Daucus carota L Planta. 1990 Jan;180(2):285-92.doi: 10.1007/BF00194009.

Knox et al.(1991). Developmentally regulated epitopes of cell surface arabinogalactan proteins and their relation to root

tissue pattern formation. Plant J. 1991 ov;1(3):317-326.doi: 10.1046/j.1365-313X.1991.t01-9-00999.x.