

Product no **AS10 659**

Rabbit anti-Goat IgG (H&L), HRP conjugated, min, cross-reactivity to human, mouse and rat serum

Product information

Immunogen | Purified Goat IgG, whole molecule

Host | Rabbit

Clonality | Polyclonal

Purity | Immunogen affinity purified rabbit IgG.

Format | Lyophilized

Quantity | 1 mg

Reconstitution | For reconstitution add 1,1 ml of sterile water, Let it stand 30 minutes at room temperature to dissolve, Prepare fresh working dilutions daily

Storage | Store lyophilized material at 2-8°C. For long time storage after reconstitution, dilute the antibody solution with glycerol to a final concentration of 50% glycerol and store as liquid at -20°C, to prevent loss of enzymatic activity. For example, if you have reconstituted 1 mg of antibody in 1,1 ml of sterile water add 1,1 ml of glycerol. Such solution will not freeze in -20°C. If you are using a 1:5000 dilution prior to diluting with glycerol, then you would need to use a 1:2500 dilution after adding glycerol. Prepare working dilution prior to use and then discard, Be sure to mix well but without foaming.

Additional information | HRP-conjugate is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 10 % (w/v) BSA, Protease/IgG free

0.1 % (v/v) of Kathon CG is used as preservative.

Application information

Recommended dilution | 1 : 10 000 -1 : 50 000 (ELISA), 1 : 500 -1 : 5000 (IHC), 1 : 10 000 -1 : 50 000 (WB)

Confirmed reactivity | Goat IgG heavy chains and light chains on all Goat immunoglobulins

Predicted reactivity | Goat IgG Heavy chains and Light chains on all Goat immunoglobulins

Not reactive in | No confirmed exceptions from predicted reactivity are currently known

Additional information | No reactivity is observed to non-immunoglobulin goat serum proteins. No reactivity is observed to serum from human, mouse and rat based on immunoelectrophoresis.

BSA and milk have to be replaced by other blocking reagents, like donkey serum or commercial formulations which are free from bovine IgG.