

Product no **AS16 3353**

Goat anti-Human IgG Fc, F(ab)'2 Fragment, DyLight® 488 conjugated, min, cross-reactivity to human IgA, IgM or IgG F(ab)'2 fragment

Product information

Immunogen | Purified Human IgG, Fc fragment

Host | Goat

Clonality | Polyclonal

Purity | Immunogen affinity purified goat IgG.

Format | Lyophilized

Quantity | 0.5 mg

Reconstitution | For reconstitution add 0.55 ml of sterile water. Let it stand 30 minutes at room temperature to dissolve. Prepare fresh working dilutions daily

Storage | Store freeze-dried powder at 2-8 °C.

Additional information | Antibody purity is > 90% based on SDS-PAGE. Small amounts of intact IgG may be present.
Affinity purified using solid phase Human IgG .
Antibody is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1% (w/v) BSA, Protease/IgG free.
0.05% (w/v) Sodium Azide is added as a preservative.

Based on IEP, this antibody reacts with: • heavy () chains on human IgG.
Based on IEP, no reactivity is observed to: • non-immunoglobulin human serum immunoglobulins • light chains on all human immunoglobulins • human IgA or IgM.

Application information

Recommended dilution | 1 : 20-1 : 2000 (Flow cyt), (IL)

Confirmed reactivity | Goat anti-Human IgG Fc, F(ab)'2 Fragment