

Product no **AS10 1018**

Goat anti-Rabbit IgG (H&L), F(ab)'2 fragment, TRITC conjugated

Product information

Host | Goat

Clonality | Polyclonal

Purity | Immunogen affinity purified IgG, F(ab)'2 fragment.

Format | Liquid

Quantity | 0.5 mg

Storage | Store at 2-8°C. Shelf life of this product is one year from date of receipt.

Additional information | This antibody reacts with the heavy chains on rabbit IgG and with the light chains on all rabbit immunoglobulins based on immunoelectrophoresis.

Minimum cross-reactivity is observed to non-immunoglobulin rabbit serum proteins based on immunoelectrophoresis.

Antibody is supplied in 10 mM sodium phosphate, 150 mM sodium chloride, pH 7.2, 1% (w/v) BSA, protease/IgG free and 0.05 % (w/v) sodium azide as preservative.

Application information

Recommended dilution | The optimal working dilution should be determined by the investigator, (ICC), immunohistochemistry (IHC) (Flow cyt)

Selected references | [Feng et al. \(2020\)](#). Intracellular expression of arginine deiminase activates the mitochondrial apoptosis pathway by inhibiting cytosolic ferritin and inducing chromatin autophagy. *BMC Cancer*. 2020 Jul 16;20(1):665.doi: 10.1186/s12885-020-07133-4.