

This product is for research use only (not for diagnostic or therapeutic use)

contact: support@agrisera.com

Agrisera AB | Box 57 | SE-91121 Vännäs | Sweden | +46 (0)935 33 000 | www.agrisera.com

Product no AS16 3349

Goat anti-Human IgG Fc, TRITC conjugated, min, cross-reactivity to human IgA + IgM, bovine, horse, mouse or rabbit serum proteins

Product information

Immunogen Purified Human IgG, Fc fragment

Host Goat

Clonality Polyclonal

Purity Immunogen affinity purified goat IgG.

Format Lyophilized

Quantity 1 mg

additing 1 in

Reconstitution For reconstitution add 1,1 ml of sterile water,Let it stand 30 minutes at room temperature to dissolve, Prepare fresh

working dilutions daily

Storage Store at 2-8°C.

Additional information Antibody purity is > 95% based on SDS-PAGE.

Affinity purified using solid phase Human IgG.

Antibody is supplied in 10 mM Sodium Phosphate, 0.15 M Sodium Chloride, pH 7.2, 1 % (w/v) BSA, Protease/lgG free.

0.05% (w/v) Sodium Azide is added as a preservative.

Based on IEP, this antibody reacts with: • heavy () chains on human IgG .

Based on IEP, no reactivity is observed to: • non-immunoglobulin human serum immunoglobulins • light chains on all

human immunoglobulins • human IgA or IgM • serum proteins from bovine, mouse, or rabbit .

Application information

Recommended dilution 1:20-1:2000 (Flow cyt), (IL)

Confirmed reactivity | Human IgG Fc