



Recombinant Cynomolgus ACVRL1 Protein, C-His-tagged

Product Information

Cat	IMP-1795
Official Symbol	ACVRL1
Product Overview	Recombinant cynomolgus ACVRL1 (XP_005570958.1) (Met1-Gln118) was expressed with a polyhistidine tag at the C-terminus.
Description	Activin A receptor, type II-like 1 (ACVRL1), also known as ALK-1 (activin receptor-like kinase 1), is an endothelial-specific type I receptor of the TGF-beta (transforming growth factor beta) receptor family of ligands. On ligand binding, a heteromeric receptor complex forms consisting of two type II and two type I transmembrane serine/threonine kinases. ACVRL1 protein is expressed in certain blood vessels of kidney, spleen, heart and intestine, serving as an important role during vascular development. Mutations in ACVRL1 gene are associated with hemorrhagic telangiectasia type 2, also known as Rendu-Osler-Weber syndrome 2 and vascular disease.
Expression System	HEK293
Species	Cynomolgus
Tag	C-His
Predicted N Terminal	Asp 22
Form	Lyophilized from sterile PBS, pH 7.4, 5 % trehalose, 5% mannitol and 0.01% Tween80.
Molecular Mass	The recombinant cynomolgus ACVRL1 consists 108 amino acids and predicts a molecular mass of 12.2 kDa.
Protein length	Met1-Gln118
Endotoxin	< 1.0 EU/μg of the protein as determined by the LAL method
Purity	> 95 % as determined by SDS-PAGE
Storage	Samples are stable for up to twelve months from date of receipt at -20 to -80 centigrade. Store it under sterile conditions at -20 to -80 centigrade. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.
Reconstitution	A hardcopy of COA with reconstitution instruction is sent along with the products. Please refer to it for detailed information.
SDS-PAGE	



Bioactivity-Cell based assay 1



Measured by its ability to inhibit BMP9-induced alkaline phosphatase production by MC3T3-E1 cells. The ED50 for this effect is typically 0.05-0.25 µg/mL.