

## **Recombinant Mouse Tnfrsf4 protein**

## **Product Information**

Cat IMP-364

Official Symbol Tnfrsf4

Product Overview Recombinant Mouse Tnfrsf4 protein(NP\_035789.1) (Met1-Pro211) was

expressed in HEK293.

**Description** OX40 (CD134) and its binding partner, OX40L (CD252), are members of

the tumor necrosis factor receptor/tumor necrosis factor superfamily, is known to break an existing state of tolerance in malignancies, leading to a reactivation of antitumor immunity. The interaction between OX40 and OX40L plays an important role in antigen-specific T-cell expansion and survival. OX40 and OX40L also regulate cytokine production from T cells, antigen-presenting cells, natural killer cells, and natural killer T cells, and

modulate cytokine receptor signaling. In line with these important

modulatory functions, OX40-OX40L interactions have been found to play a central role in the development of multiple inflammatory and autoimmune diseases, making them attractive candidates for intervention in the clinic. Conversely, stimulating OX40 has shown it to be a candidate for

therapeutic immunization strategies for cancer and infectious disease.

Expression System HEK293

**Species** Mouse

Tag N/A

Predicted N Terminal Val 20

**Form** Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose,

mannitol and 0.01% Tween80 are added as protectants before

lyophilization.

Molecular Mass The recombinant mouse Tnfrsf4 consists 199 amino acids and predicts a

molecular mass of 22.1 kDa.

Protein length Met1-Pro211

**Endotoxin** < 1.0 EU per μg protein as determined by the LAL method.

**Purity** > 95 % as determined by SDS-PAGE.

Storage Samples are stable for up to twelve months from date of receipt at -20°C to

-80°C. Store it under sterile conditions at -20°C to -80°C. It is

recommended that the protein be aliquoted for optimal storage. Avoid

repeated freeze-thaw cycles.

**Reconstitution** It is recommended that sterile water be added to the vial to prepare a stock

solution of 0.2 ug/ul. Centrifuge the vial at 4°C before opening to recover



the entire contents.