

## Streptavidin Coated Plates, Clear, 96-Well

**Catalog Number :** SP-11

**Pack Size:** 1 plate / 5 plates

### Specifications

Table 1. plate details

Items	Specifications
Material	Polystyrene
Color	Clear
Plate Blocking:	2% BSA Blocking Buffer
Formulations	Clear, 96-well plates, coated with 100uL of streptavidin tetramer and blocked with 200uL of 2% BSA Blocking Buffer
Detection Method	Colorimetric
Capacity	~5pmol biotin/well
CV% of plates/wells	< 10%
Type	Detection Plate, Immunoassay, ELISA

### Shipping and Storage

Upon receipt store plates at 4°C in unopened pouches. Once opened, place unused plates in a resealable bag with desiccant and store at 4°C. This plate is supplied and shipped with blue ice.

### Product description

The Streptavidin Coated Plates, Clear, 96-Well is pre-coated with Streptavidin tetramer protein and blocked with BSA, it is a ready-to-use polystyrene plate, which can be used for binding biotinylated proteins and antibodies, or probes for ELISA and other target-specific assays. The recombinant Streptavidin is tetramer protein expressed in E. coli designed for immobilization applications.

### Applications

This Streptavidin Coated Plate is intended for Immunoassay and ELISA.

**IMPORTANT:** Please carefully read this manual before performing your experiment.

**For Research Use Only. Not For Use In Diagnostic Or Therapeutic Procedures**

**US and Canada:**

**Tel:** +1 800-810-0816

**Web:** <http://www.acrobiosystems.com>

**Asia and Pacific:**

**Tel:** +86 400-682-2521

**E-mail:** [order@acrobiosystems.com](mailto:order@acrobiosystems.com)

## **Assay Principles**

Streptavidin (SA) has an extraordinarily high affinity for biotin with a dissociation constant (Kd) on the order of  $10^{-14}$  mol/L, the Biotinylated molecules can bind to the SA irreversibly. Streptavidin has an isoelectric point of 5 to 6, resulting in low nonspecific interactions. The Streptavidin Coated Plates we provide are easy to use and widely available for applications.

## **Example ELISA Procedure**

### **Materials and Reagents Preparation**

Before starting the ELISA Assay, we should prepare the all reagents and materials required in the experiment. You can prepare these reagents by following operations.

**Wash Buffer:** PBS or TBS with 0.05% (v/v) Tween-20 (usually at pH7.4), 500 mL is sufficient for 96 tests. The pH of Buffer system can be adjusted according to your experiment.

**Dilution Buffer:** Wash Buffer with 0.5% (w/v) bovine serum albumin (BSA) (i.e. Jackson, Catalog#. 001-000-162), 50 mL is sufficient for 96 tests.

**Substrate Dilution Buffer:** 50 mM disodium hydrogen phosphate ( $\text{Na}_2\text{HPO}_4$ ) and 25 mM citric acid, adjust pH to 5.5 with 1 M Sodium hydroxide (NaOH), 25 mL is sufficient for 96 tests.

**Substrate Stock Solution:** 20 mg/mL TMB (*Sigma-Aldrich, Catalog # 860336*) in Dimethyl sulfoxide (*Sigma-Aldrich, Catalog # D8418*), 1 mL is sufficient for 96 tests. **Protect from light.**

### **TMB Substrate Working Solution**

For **each plate** dilute 125  $\mu\text{L}$  substrate stock solution in 25 mL substrate dilution buffer and add 20  $\mu\text{L}$  5%  $\text{H}_2\text{O}_2$  (pipette 10  $\mu\text{L}$  30%  $\text{H}_2\text{O}_2$  into 50  $\mu\text{L}$  distilled water), mix well.

### **Notes:**

- 1) The TMB Substrate Working Solution should be freshly prepared and used within 15 minutes.
- 2) If you choose to use other commercially available ready-to-use TMB substrate solutions, you should follow the manufacturer's instruction.

**Stop Solution:** 1 M sulfuric acid (aqueous), 6 mL is sufficient for 96 tests.

**Microplate sealing film** (*Sigma-Aldrich, Catalog # Z724742*)

### **Pipettes and pipette tips**

**UV/Vis microplate spectrophotometer** (absorbance 450 nm, correction wavelength set to 630 nm).

---

## Recommended Protocol

### 1. Preparation

Reconstitute and store all reagents as recommended.

### 2. Washing

Add 300  $\mu$ L of Wash buffer to each well, gently tap the plate for 1 minute, remove any remaining Wash Buffer by aspirating or decanting, invert the plate and blot it against paper towels. Repeat the wash step above for three times.

### 3. Add biotinylated protein or antibodies

- 1) Dilute Biotinylated protein or antibodies to a concentration you want (usually 1~10  $\mu$ g/mL) with Dilution Buffer to make Biotinylated molecule working solution.
- 2) Add 100  $\mu$ L Biotinylated molecule to each well and incubate at 37°C or RT for 1 hour.
- 3) For the "Blank" wells, please add 10  $\mu$ L Biotinylated molecule working solution.
- 4) For Non - specific of the sample wells, please add 100  $\mu$ L Dilution Buffer.

### 4. Washing

Remove the remaining solution by aspiration, add 300  $\mu$ L of Wash buffer to each well, gently tap the plate for 1 minute, remove any remaining Wash Buffer by aspirating or decanting, invert the plate and blot it against paper towels. Repeat the wash step above **for three times**.

### 5. Add Samples

- 1) Make series dilution of the samples as appropriate with Dilution Buffer.
- 2) Add 100  $\mu$ L of the serial dilution of sample to each well, incubate at 37°C or RT for 1 hour.

### 6. Washing

Repeat step 4.

### 7. Add primary antibody

- 1) Dilute primary antibody to an appropriate concentration with Dilution Buffer.
- 2) For all wells, add 100  $\mu$ L of diluted primary antibody, and incubate at 37°C or RT for 1 hour.

### 8. Washing

Repeat step 4.

### 9. Add enzyme-labeled secondary antibody

- 1) Dilute secondary antibody to an appropriate concentration with Dilution Buffer.
- 2) For all wells, add 100  $\mu$ L of diluted secondary antibody, and incubate at 37°C or RT for 1 hour, avoid light.

### 10. Washing

Repeat step 4.

### 11. TMB Substrate Reaction

Add 200  $\mu$ L TMB Substrate Working Solution to each well. Seal the plate with microplate sealing film and incubate at 37°C or RT for 20 minutes, avoid light.

### 12. Termination

Add 50  $\mu$ L **Stop Solution** to each well, and tap the plate gently for 3 minutes to allow thorough mixing.

*Note: the color in the wells should change from blue to yellow.*

### 13. Data Recording

Read the absorbance at 450 nm using UV/Vis microplate spectrophotometer.


*Note: the plate may be read at 600 nm without adding 1 M sulfuric acid, but the Signal-to-Background ratio*

may be reduced.

### Example Data


#### 1. Binding Assay between CD19 and Anti-CD19 antibody on SA Plate

Immobilized Biotinylated Human CD19 (20-291), His,Avitag (Cat. No. CD9-H82E9) at 1 µg/mL (100 µL/well) on Streptavidin Coated Plates, Clear, 96-Well (Cat. No. SP-11), can bind Anti-FMC63 antibody with a linear range of 0.1-3 ng/mL (QC tested).




#### 2. Binding Assay between PD1 and PDL1 on SA Plate

Immobilized Biotinylated Human PD-1, Fc, Avi tag, His Tag (Cat. No. PD1-H82F4) at 1 µg/mL (100 µL/well) on Streptavidin Coated Plates, Clear, 96-Well (Cat. No. SP-11), can bind Human PD-L1, mFc Tag with a linear range of 0.4-6.25 ng/mL (QC tested).




### 3. Binding Assay between CD20 and Rituximab on SA Plate

Immobilized Biotinylated Human CD20 Full Length, His, Avi tag (HEK293) (Cat. No. CD0-H82E5) at 1  $\mu\text{g/mL}$  (100  $\mu\text{L/well}$ ) on Streptavidin Coated Plates, Clear, 96-Well (Cat. No. SP-11), can bind Rituximab with a linear range of 78-625 ng/mL (QC tested).



### 4. Binding Assay between FcRn and Ipilimumab on SA Plate

Immobilized Biotinylated Human FCGRT&B2M Heterodimer Protein, His, Avi tag (Cat. No. FCM-H82W7) at 1  $\mu\text{g/mL}$  (100  $\mu\text{L/well}$ ) on Streptavidin Coated Plates, Clear, 96-Well (Cat. No. SP-11), can bind Ipilimumab with a linear range of 78-1250 ng/mL (QC tested).



## TROUBLESHOOTING GUIDE

Problem	Possible Cause	Solutions
Signal of positive control is weak or abnormal	Incorrect storage of plate	✧ The plate should be store plates at 4°C, once you open the package, get the amount you need and keep the rest airtight.
	Detection Antibody is outdated or no prepared the working solution immediately before use	✧ The working solution should be prepared immediately before use and should not be stored.
	Errors in instrument settings	✧ Please check instrument setting.
	Substrate Stock Solution is outdated; Incubation temperature is incorrect; Incubation time is not sufficient; Repeated freeze-thaw cycles;	✧ Make sure the <b>Substrate Stock Solution</b> is working. ✧ Use proper incubation time and temperature.
	Pipetting errors	✧ Make sure that the pipette is calibrated and working properly.
High background	Serum samples	✧ If you want test serum samples, the BSA Blocking plate is not suitable for this purpose. <b>We specially developed the Streptavidin Coated Plates, Clear,96-Well (For Serological Testing) (Cat.No.SP-13) for serological testing.</b>
	Sample solvent contains inhibiting factors	✧ Run a negative control assay with the solvent alone. ✧ Maintain DMSO level at <1%. Increase protein incubation time.
	Contamination	✧ Make sure buffers and samples are prepared, used and stored correctly.
	The <b>TMB Substrate Working Solution</b> is not fresh	✧ <b>TMB Substrate Working Solution</b> must be used within 15 minutes after preparation.
Colorimetric signal is erratic	Inconsistent pipetting or dilution methods	✧ Make sure pipettors are functioning properly and use a multichannel pipettor if possible. ✧ Use master mixes to minimize errors. ✧ Run duplicates for all tests.
	<b>TMB Substrate Working Solution</b> is not completely mixed with the reaction solution	✧ Make sure that <b>TMB Substrate Working Solution</b> is adequately mixed with the reaction solution.
	Bubbles in the wells	✧ Tap plate gently to disperse bubbles.
	Signal is too high	✧ The concentration of the samples should be adjusted to achieve optimal reading. ✧ Decrease colorimetric HRP substrate incubation time.
Inadequate color development	Incomplete removal of residual buffers during previous steps	✧ Wells should appear dry after aspiration.
	Problems with conjugate or color reagents	✧ Color should appear immediately after the reagent is added. Make sure no contamination or residual buffers in the wells before you start the color development process.