



## Synonym

HLA-A\*0201 & B2M & EBV EBNA (FMVFLQTHI)

## Source

Biotinylated Human HLA-A\*02:01&B2M&EBV EBNA (FMVFLQTHI) Complex Protein(HLA-H82E8) is expressed from human 293 cells (HEK293). It contains AA Gly 25 - Ile 308 (HLA-A\*02:01) & Ile 21 - Met 119 (B2M) & FMVFLQTHI peptide (Accession # [AAA59606.1](#) (HLA-A\*02:01) & [P61769-1](#) (B2M) & FMVFLQTHI).

Predicted N-terminus: Gly 25 & Ile 21

## Molecular Characterization

Biotinylated Human HLA-A\*02:01&B2M&EBV EBNA (FMVFLQTHI) Complex Protein is produced by co-expression of HLA and B2M loaded with EBV EBNA peptide.

This protein carries a polyhistidine tag at the C-terminus, followed by an Avi tag (Avitag™).

The protein has a calculated MW of 36.3 kDa and 11.7 kDa. The protein migrates as 41-43 kDa and 10 kDa when calibrated against [Star Ribbon Pre-stained Protein Marker](#) under reducing (R) condition (SDS-PAGE) due to glycosylation.

## Labeling

*Biotinylation of this product is performed using Avitag™ technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.*


## Protein Ratio

Passed as determined by the HABA assay / binding ELISA.

## Endotoxin

Less than 1.0 EU per µg by the LAL method.

## SDS-PAGE



Biotinylated Human HLA-A\*02:01&B2M&EBV EBNA (FMVFLQTHI) Complex Protein on SDS-PAGE under reducing (R) condition. The gel was

## Purity

>90% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

## Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.

Contact us for customized product form or formulation.

## Reconstitution

Please see Certificate of Analysis for specific instructions.

*For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.*

## Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

*Please avoid repeated freeze-thaw cycles.*

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70°C for 3 months under sterile conditions after reconstitution.

## SEC-MALS



The purity of Biotinylated Human HLA-A\*02:01&B2M&EBV EBNA (FMVFLQTHI) Complex Protein (Cat. No. HLA-H82E8) is more than 90%

**Discounts, Gifts,  
and more!**



» [www.acrobiosystems.com](http://www.acrobiosystems.com)

# Biotinylated Human HLA-A\*02:01&B2M&EBV EBNA (FMVFLQTHI) Complex Protein (Monomer, MALS verified)

Catalog # HLA-H82E8



BIOSYSTEMS  
**Acro**  
Surprise Inside!


stained with Coomassie Blue. The purity of the protein is greater than 90% (With [Star Ribbon Pre-stained Protein Marker](#)).

and the molecular weight of this protein is around 45-60 kDa verified by SEC-MALS.


[Report](#)

## Bioactivity-ELISA

**Biotinylated Human HLA-A\*02:01&B2M&EBV EBNA (FMVFLQTHI) Complex Protein ELISA**  
0.1 µg of Anti-HLA class I Antibody, Human IgG1 (W6/32) per well




**Biotinylated Human HLA-A\*02:01&B2M&EBV EBNA (FMVFLQTHI) Complex Protein ELISA**  
0.1 µg of Biotinylated Human HLA-A\*02:01&B2M&EBV EBNA (FMVFLQTHI) Complex Protein per well



Immobilized Anti-HLA class I Antibody, Human IgG1 (W6/32) at 1 µg/mL (100 µL/well) can bind Biotinylated Human HLA-A\*02:01&B2M&EBV EBNA (FMVFLQTHI) Complex Protein (Cat. No. HLA-H82E8) with a linear range of 0.5-8 ng/mL (QC tested).

Immobilized Biotinylated Human HLA-A\*02:01&B2M&EBV EBNA (FMVFLQTHI) Complex Protein (Cat. No. HLA-H82E8) at 1 µg/mL (100 µL/well) on streptavidin (Cat. No. STN-N5116) precoated (0.5 µg/well) plate can bind Anti-HLA class I Antibody, Human IgG1 (W6/32) with a linear range of 0.2-3 ng/mL (Routinely tested).

**Biotinylated Human HLA-A\*02:01&B2M&EBV EBNA (FMVFLQTHI) Complex Protein ELISA**  
0.1 µg of Biotinylated Human HLA-A\*02:01&B2M&EBV EBNA (FMVFLQTHI) Complex Protein per well



Immobilized Biotinylated Human HLA-A\*02:01&B2M&EBV EBNA (FMVFLQTHI) Complex Protein (Cat. No. HLA-H82E8) at 1 µg/mL (100 µL/well) on streptavidin (Cat. No. STN-N5116) precoated (0.5 µg/well) plate can bind Anti-B2M Antibody, Human IgG1 with a linear range of 0.2-3 ng/mL (Routinely tested).

## Clinical and Translational Updates

Discounts, Gifts,  
and more!



» [www.acrobiosystems.com](http://www.acrobiosystems.com)