

Villin Antibody / VIL1 [clone VIL1/7378R] (V4949)

| Catalog No. | Formulation | Size |
|----------------|---|--------|
| V4949-100UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 100 ug |
| V4949-20UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 20 ug |
| V4949SAF-100UG | 1 mg/ml in 1X PBS; BSA free, sodium azide free | 100 ug |


Recombinant RABBIT MONOCLONAL

Bulk quote request

| Availability | 1-3 business days |
|--------------------|--|
| Species Reactivity | Human |
| Format | Purified |
| Clonality | Recombinant Rabbit Monoclonal |
| Isotype | Rabbit IgG, kappa |
| Clone Name | VIL1/7378R |
| Purity | Protein A/G affinity |
| UniProt | P09327 |
| Localization | Cytoplasm, Cell Surface |
| Applications | Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT |
| Limitations | This Villin antibody is available for research use only. |



IHC staining of FFPE human colon adenocarcinoma with Villin antibody (clone VIL1/7378R). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human colon adenocarcinoma with Villin antibody (clone VIL1/7378R). Inset: PBS used in place of primary Ab (secondary Ab negative control). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.

Description

Recognizes a protein of 95kDa, which is identified as villin. It is a major constituent in the microvilli, which compose the brush border of epithelial cells forming absorptive surfaces of the intestinal and renal proximal tubular epithelia. Anti-Villin labels the brush border area in the gastrointestinal mucosal epithelium and urogenital tract. Among neoplasms, villin is predominantly expressed in tumors of colorectal origin. Antibody to villin is useful in identifying malignant cells from primary and metastatic colorectal carcinomas. This antibody also labels Merkel cells of the skin.

Application Notes

Optimal dilution of the Villin antibody should be determined by the researcher.

Immunogen

A recombinant partial protein sequence (within amino acids 600-700) from the human protein was used as the immunogen for the Villin antibody.

Storage

Aliquot the Villin antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.

Ordering:Phone:858.663.9055 | Fax:1.267.821.0800 | Email:info@nsjbio.com

Copyright © NSJ Bioreagents. All rights reserved