NSJ BIOREAGENTS

MRP4 Antibody / Multidrug resistance-associated protein 4 / ABCC4 [clone ABCC4/9179] (V5382)

Catalog No.	Formulation	Size
V5382-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V5382-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V5382SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request

Availability	1-3 business days	
Species Reactivity	Human	
Format	Purified	
Clonality	Monoclonal (mouse origin)	
Isotype	Mouse IgG2b, kappa	
Clone Name	ABCC4/9179	
Purity	Protein A/G affinity	
UniProt	O15439	
Localization	Membrane	
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml	
Limitations	This MRP4 antibody is available for research use only.	



IHC staining of FFPE human prostate cancer tissue with MRP4 antibody (clone ABCC4/9179). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.

kDa R NR	SDS-PAGE analysis of purified, BSA-free MRP4 antibody (clone ABCC4/9179) as confirmation of integrity and purity.
250	
150 — 🛩	
100	
75	
50 =	
37	
25 2ug loadir	g
15 NR=Non-	
10 — Bereduced	

Description

The two members of the large family of ABC transporters known to confer multidrug resistance in human cancer cells are the Mdr-1 P-glycoprotein and the multidrug-resistance protein MRP1. MRP1 is an integral membrane protein that contains an MDR-like core, an N-terminal membrane-bound region and a cytoplasmic linker, and it is expressed in various cerebral cells, as well as in lung, testis and peripheral blood. The MRP gene family also includes MRP2, which is alternatively designated cMOAT (for canalicular multispecific organic anion transporter), and MRP3, which are both conjugate export pumps expressed predominantly in hepatocytes. MRP2 localizes exclusively to the apical membrane and is constitutively expressed at a high level in normal liver cells. Conversely, MRP3 localizes to the basolateral membrane where it also mediates the transport of the organic anion S-(2,4-dinitrophenyl-) glutathione toward the basolateral side of the membrane. MRP3 is normally expressed at comparatively lower levels than MRP2 and increases only when secretion across the apical membrane by MRP2 is impaired. MRP6 is highly expressed in liver and kidney, whereas MRP4 and MRP5 are detected in various tissues, yet at much lower levels of expression.

Application Notes

Optimal dilution of the MRP4 antibody should be determined by the researcher.

Immunogen

A recombinant fragment (within amino acids 1-200) of human ABCC4 protein was used as the immunogen for the MRP4 antibody.

Storage

Aliquot the MRP4 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.

Ordering:Phone:858.663.9055 | Fax:1.267.821.0800 | Email:info@nsjbio.com

Copyright © NSJ Bioreagents. All rights reserved