


IL17D Antibody (F44066)

| Catalog No. | Formulation | Size |
|---------------|--|---------|
| F44066-0.4ML | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.4 ml |
| F44066-0.08ML | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.08 ml |

[Bulk quote request](#)

| | |
|---------------------------|---|
| Availability | 1-3 business days |
| Species Reactivity | Human |
| Format | Antigen affinity purified |
| Clonality | Polyclonal (rabbit origin) |
| Isotype | Rabbit Ig |
| Purity | Antigen affinity |
| UniProt | Q8TAD2 |
| Applications | Western blot : 1:1000 |
| Limitations | This IL17D antibody is available for research use only. |



IL17D antibody western blot analysis in NCI-H460 lysate.

Description

The protein encoded by this gene is a cytokine that shares the sequence similarity with IL17. The treatment of endothelial cells with this cytokine has been shown to stimulate the production of other cytokines including IL6, IL8 and CSF2/ GM-CSF. The increased expression of IL8 induced by this cytokine was found to be NF-kappa B-dependent.

Application Notes

Titration of the IL17D antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

A portion of amino acids 155-183 from the human protein was used as the immunogen for this IL17D antibody.

Storage

Aliquot the IL17D antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.