


NPM1 Antibody (F48247)


| Catalog No. | Formulation | Size |
|---------------|--|---------|
| F48247-0.4ML | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.4 ml |
| F48247-0.08ML | In 1X PBS, pH 7.4, with 0.09% sodium azide | 0.08 ml |

[Bulk quote request](#)


| | |
|-----------------------------|---|
| Availability | 1-3 business days |
| Species Reactivity | Human |
| Predicted Reactivity | Mouse, Rat |
| Format | Purified |
| Clonality | Polyclonal (rabbit origin) |
| Isotype | Rabbit Ig |
| Purity | Purified |
| UniProt | P06748 |
| Applications | Western blot : 1:1000 Immunofluorescence : 1:10-1:50 IHC (Paraffin) : 1:10-1:50 Flow Cytometry : 1:10-1:50 |
| Limitations | This NPM1 antibody is available for research use only. |




IHC analysis of FFPE human lymph tissue stained with NPM1 antibody



Fluorescent image of A549 cell stained with NPM1 antibody. Alexa Fluor 488 secondary was used (green). NPM1 immunoreactivity is localized to the nucleus and nucleolus.



NPM1 antibody western blot analysis in HeLa lysate. Expected/observed molecular weight: ~38kDa.



NPM1 antibody flow cytometric analysis of HeLa cells (right histogram) compared to a [negative control](#) (left histogram). FITC-conjugated donkey-anti-rabbit secondary Ab was used for the analysis.

Description

Nucleophosmin/NPM1 is a ubiquitously expressed nucleolar protein that shuttles between the nucleus and cytoplasm. It is implicated in multiple functions, including ribosomal protein assembly and transport, control of centrosome duplication, and regulation of the tumor suppressor ARF. NPM1 mutations that relocalize NPM1 from the nucleus into the cytoplasm are associated with development of acute myeloid leukemia.

Application Notes

Titration of the NPM1 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

A portion of amino acids 7-33 from the human protein was used as the immunogen for this NPM1 antibody.

Storage

Aliquot the NPM1 antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.

