


Beta-TrCP Antibody / BTRC (F54622)

Catalog No.	Formulation	Size
F54622-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F54622-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml

[Bulk quote request](#)


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit Ig
Purity	Antigen affinity purified
UniProt	Q9Y297
Localization	Cytoplasmic, nuclear
Applications	Immunofluorescence : 1:25 Immunohistochemistry (FFPE) : 1:25 Western blot : 1:500-1:2000
Limitations	This Beta-TrCP antibody is available for research use only.




Western blot testing of human 1) HepG2 and 2) HEK293 cell lysate with Beta-TrCP antibody. Predicted molecular weight ~69 kDa.

kDa
95
72
55
36
28
17

Western blot testing of human ZR-75-1 cell lysate with Beta-TrCP antibody. Predicted molecular weight ~69 kDa.



Immunofluorescent staining of human ZR-75-1 cells with Beta-TrCP antibody (green), DAPI nuclear stain (blue) and anti-Actin (red).



IHC testing of FFPE human brain tissue with Beta-TrCP antibody. HIER: steam section in pH6 citrate buffer for 20 min and allow to cool prior to staining.

Description

This gene encodes a member of the F-box protein family which is characterized by an approximately 40 amino acid motif, the F-box. The F-box proteins constitute one of the four subunits of ubiquitin protein ligase complex called SCFs (SKP1-cullin-F-box), which function in phosphorylation-dependent ubiquitination. The F-box proteins are divided into 3 classes: Fbws containing WD-40 domains, Fbls containing leucine-rich repeats, and Fbxs containing either different protein-protein interaction modules or no recognizable motifs. The protein encoded by this gene belongs to the Fbws class; in addition to an F-box, this protein contains multiple WD-40 repeats. This protein is homologous to *Xenopus* bTrCP1, yeast Met30, *Neurospora* Scon2 and *Drosophila* Slimb proteins. It interacts with HIV-1 Vpu and connects CD4 to the proteolytic machinery. It also associates specifically with phosphorylated I κ B α and beta-catenin destruction motifs, probably functioning in multiple transcriptional programs by activating the NF- κ B pathway and inhibiting the beta-catenin pathway.

Application Notes

The stated application concentrations are suggested starting points. Titration of the Beta-TrCP antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

A portion of amino acids 17-52 from the human protein was used as the immunogen for the Beta-TrCP antibody.

Storage

Aliquot the Beta-TrCP antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.