


Mitochondrial amidoxime reducing component 2 Antibody / mARC2 / MTARC2 (F54991)


Catalog No.	Formulation	Size
F54991-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F54991-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml

[Bulk quote request](#)


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit Ig
Purity	Antigen affinity purified
UniProt	Q969Z3
Localization	Cytoplasmic
Applications	Western blot : 1:500-1:1000 Flow cytometry : 1:10-1:50 (1x10e6 cells) Immunohistochemistry (FFPE) : 1:50-1:100
Limitations	This Mitochondrial amidoxime reducing component 2 antibody is available for research use only.




IHC testing of FFPE human lung carcinoma tissue with Mitochondrial amidoxime reducing component 2 antibody. HIER: steam section in pH6 citrate buffer for 20 min and allow to cool prior to staining.



Western blot testing of human HeLa cell lysate with Mitochondrial amidoxime reducing component 2 antibody. Predicted molecular weight ~38 kDa.



Western blot testing of 1) non-transfected and 2) transfected 293 cell lysate with Mitochondrial amidoxime reducing component 2 antibody.



Flow cytometry testing of human HeLa cells with Mitochondrial amidoxime reducing component 2 antibody; Blue=isotype control, Green= Mitochondrial amidoxime reducing component 2 antibody.

Description

Catalytic component of the benzamidoxime prodrug-converting complex, a complex required to reduce N-hydroxylated structures, such as benzamidoxime prodrug. Benzamidoxime is an amidine prodrug produced by N-hydroxylation which is used to enhance bioavailability and increase intestinal absorption. It is then reduced into benzamidine, its active amidine, by the benzamidoxime prodrug-converting complex.

Application Notes

The stated application concentrations are suggested starting points. Titration of the Mitochondrial amidoxime reducing component 2 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

A portion of amino acids 270-300 from the human protein was used as the immunogen for the Mitochondrial amidoxime reducing component 2 antibody.

Storage

Aliquot the Mitochondrial amidoxime reducing component 2 antibody and store frozen at -20oC or colder. Avoid repeated

freeze-thaw cycles.

Ordering:Phone:858.663.9055 | Fax:1.267.821.0800 | Email:info@nsjbio.com

Copyright © NSJ Bioreagents. All rights reserved