


## MPI Antibody / Mannose Phosphate Isomerase (R32582)


Catalog No.	Formulation	Size
R32582	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

**Bulk quote request**


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human, Mouse, Rat
<b>Format</b>	Antigen affinity purified
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit IgG
<b>Purity</b>	Antigen affinity
<b>Buffer</b>	Lyophilized from 1X PBS with 2% Trehalose and 0.025% sodium azide
<b>UniProt</b>	P34949
<b>Localization</b>	Cytoplasmic
<b>Applications</b>	Western blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml Flow cytometry : 1-3ug/10 <sup>6</sup> cells
<b>Limitations</b>	This MPI antibody is available for research use only.




Western blot testing of 1) rat testis and 2) human HeLa lysate with MPI antibody at 0.5ug/ml. Predicted molecular weight ~47 kDa.




IHC staining of FFPE human intestinal cancer with MPI antibody at 1ug/ml. HIER: boil tissue sections in pH6, 10mM citrate buffer, for 10-20 min followed by cooling at RT for 20 min.




IHC staining of FFPE human lung cancer with MPI antibody at 1ug/ml. HIER: boil tissue sections in pH6, 10mM citrate buffer, for 10-20 min followed by cooling at RT for 20 min.




IHC staining of FFPE rat brain with MPI antibody at 1ug/ml. HIER: boil tissue sections in pH6, 10mM citrate buffer, for 10-20 min followed by cooling at RT for 20 min.



Western blot testing of 1) rat ovary and 2) mouse lung lysate with MPI antibody at 0.5ug/ml. Predicted molecular weight ~47 kDa.



Flow cytometry testing of human A549 cells with MPI antibody at 1ug/10<sup>6</sup> cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= MPI antibody.



Western blot testing of 1) human HeLa, 2) human Caco-2, 3) human HEK293, 4) rat brain, 5) rat lung, 6) rat heart, 7) mouse brain and 8) mouse lung tissue lysate with MPI antibody at 0.5ug/ml. Predicted molecular weight ~47 kDa.

## Description

Mannose-6 phosphate isomerase (MPI), also called Phosphomannose isomerase (PMI), is an enzyme which facilitates the interconversion of fructose 6-phosphate (F6P) and mannose-6-phosphate (M6P). It also plays a critical role in maintaining the supply of D-mannose derivatives, which are required for most glycosylation reactions. Mutations in the MPI gene were found in patients with carbohydrate-deficient glycoprotein syndrome, type Ib. Alternative splicing results in multiple transcript variants. This MPI gene is mapped to 15q24.1.

## Application Notes

Optimal dilution of the MPI antibody should be determined by the researcher.

## Immunogen

A human recombinant protein corresponding to amino acids A2-K99 was used as the immunogen for the MPI antibody.

## Storage

After reconstitution, the MPI antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.