Fibrinogen alpha chain Antibody / FGA (R32595)

| Catalog No. | Formulation | Size |
|-------------|---|--------|
| R32595 | 0.5mg/ml if reconstituted with 0.2ml sterile DI water | 100 ug |

Bulk quote request

| Availability | 1-3 business days |
|----------------------|--|
| Species Reactivity | Human, Rat |
| Predicted Reactivity | Mouse |
| Format | Antigen affinity purified |
| Clonality | Polyclonal (rabbit origin) |
| Isotype | Rabbit IgG |
| Purity | Antigen affinity |
| Buffer | Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide |
| UniProt | P02671 |
| Localization | Secreted |
| Applications | Western blot : 0.5-1ug/ml |
| Limitations | This Fibrinogen alpha chain antibody is available for research use only. |

| kDa 200 116 97 | Western blot testing of 1) rat skeletal muscle and 2) human HepG2 cell lysate with Fibrinogen alpha chain antibody at 0.5ug/ml. Predicted molecular weight ~95 kDa (alpha-E isoform). | |
|-------------------------|---|--|
| 66 | | |
| 44 | | |
| 31 | | |
| 22 | | |
| 14 6- 1 2 | | |

Description

Fibrinogen alpha chain is a protein that in humans is encoded by the FGA gene. This gene encodes the alpha subunit of the

coagulation factor fibrinogen, which is a component of the blood clot. Following vascular injury, the encoded preproprotein is proteolytically processed by thrombin during the conversion of fibrinogen to fibrin. Mutations in this gene lead to several disorders, including dysfibrinogenemia, hypofibrinogenemia, afibrinogenemia and renal amyloidosis. Alternative splicing results in multiple transcript variants, at least one of which encodes an isoform that undergoes proteolytic processing.

Application Notes

Optimal dilution of the Fibrinogen alpha chain antibody should be determined by the researcher.

Immunogen

Amino acids 687-727 (RTWQDYKRGFGSLNDEGEGEFWLGNDYLHLLTQRGSVLRVE-human) were used as the immunogen for the Fibrinogen alpha chain antibody. This sequence is specific to the alpha-E isoform and not found in the alpha isoform.

Storage

After reconstitution, the Fibrinogen alpha chain antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.

Ordering:Phone:858.663.9055 | Fax:1.267.821.0800 | Email:info@nsjbio.com

Copyright © NSJ Bioreagents. All rights reserved