


Prothrombin Antibody (R32993)


Catalog No.	Formulation	Size
R32993	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human
Format	Antigen affinity purified
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose and 0.025% sodium azide
UniProt	P00734
Localization	Secreted
Applications	Western Blot : 0.5-1ug/ml IHC (FFPE) : 1-2ug/ml Direct ELISA : 0.1-0.5ug/ml
Limitations	This Prothrombin antibody is available for research use only.



IHC testing of FFPE human colon cancer tissue with Prothrombin antibody at 1ug/ml. Required HIER: steam section in pH6 citrate buffer for 20 min and allow to cool prior to testing.



IHC testing of FFPE human lung cancer tissue with Prothrombin antibody at 1ug/ml. Required HIER: steam section in pH6 citrate buffer for 20 min and allow to cool prior to testing.



Western blot testing of human plasma lysate with Prothrombin antibody at 0.5ug/ml. Predicted molecular weight ~70 kDa, but may be observed at higher molecular weights due to glycosylation.

Description

F2 (Coagulation Factor II), also known as prothrombin, is a serine protease that in humans is encoded by the F2 gene. This gene for human prothrombin (F2) was assigned to chromosome 11p11-q12 by analysis of a panel of somatic cell hybrid DNAs and by in situ hybridization, using both cDNA and genomic probes. The activated thrombin enzyme plays an important role in hemostasis and thrombosis: it converts fibrinogen to fibrin for blood clot formation, stimulates platelet aggregation, and activates coagulation factors V, VIII (F8), and XIII (F13A1). Thrombin also inhibits coagulation by activating protein C.

Application Notes

Optimal dilution of the Prothrombin antibody should be determined by the researcher.

Immunogen

A recombinant human partial protein corresponding to amino acids Y67-R124 was used as the immunogen for the Prothrombin antibody.

Storage

After reconstitution, the Prothrombin antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.