


GFAP Antibody [clone 3F2] (RQ6032)


Catalog No.	Formulation	Size
RQ6032	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1
Clone Name	3F2
Purity	Affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose and 0.025% sodium azide
UniProt	P14136
Localization	Cytoplasmic
Applications	Western blot : 0.5-1ug/ml Immunohistochemistry : 1-2ug/ml Immunofluorescence : 5ug/ml
Limitations	This GFAP antibody is available for research use only.




IHC staining of FFPE rat brain with GFAP antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human glioma with GFAP antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Western blot testing of 1) rat brain and 2) mouse brain lysate with GFAP antibody. Predicted molecular weight ~50 kDa.



Immunofluorescent staining of FFPE rat brain tissue with GFAP antibody (red) and DAPI nuclear stain (blue). HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.

Description

Glial fibrillary acidic protein (GFAP) is a protein that is encoded by the GFAP gene in humans. It is an intermediate filament (IF) protein that is expressed by numerous cell types of the central nervous system (CNS) including astrocytes, and ependymal cells. It is mapped to 17q21.31. GFAP is closely related to its non-epithelial family members, vimentin, desmin, and peripherin, which are all involved in the structure and function of the cell's cytoskeleton. GFAP is thought to help to maintain astrocyte mechanical strength, as well as the shape of cells. This gene has been shown to play a role in mitosis by adjusting the filament network present in the cell. GFAP is necessary for many critical roles in the CNS. What's more, GFAP also plays a role in astrocyte-neuron interactions as well as cell-cell communication.

Application Notes

Optimal dilution of the GFAP antibody should be determined by the researcher.

Immunogen

Recombinant human protein (amino acids Q93-M432) was used as the immunogen for the GFAP antibody.

Storage

After reconstitution, the GFAP antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.

