GM-CSF Antibody [clone BVD2-21C11] (V2102)

Catalog No.	Formulation	Size
V2102-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2102-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2102SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug



Bulk quote request

Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (rat origin)
Isotype	Rat IgG2a, kappa
Clone Name	BVD2-21C11
Purity	Protein G affinity chromatography
Buffer	1X PBS, pH 7.4
Gene ID	1437
Localization	Secreted (extracellular)
Applications	Neutralization Studies (order BSA/sodium azide-free format) : Immunoprecipitation : 0.5-1ug/500ug protein lysate Flow cytometry : 0.5-1ug/10^6 cells Immunofluorescence : 0.5-1ug/ml Western blot : 0.5-1ug/ml Immunohistochemistry (Frozen) : 0.5-1ug/ml for 30 min at RT
Limitations	This GM-CSF antibody is available for research use only.

Description

GM-CSF (Granulocyte/macrophage - Colony-stimulating factor) is a hematopoietic factor that is produced by activated Tcells, B-cells, mast cells, macrophages, fibroblasts, and endothelial cells. In addition to supporting colony formation of granulocyte/macrophage progenitors, GM-CSF is a growth factor for erythroid, megakaryocyte, and eosinophil progenitors.

Application Notes

The concentration stated for each application is a general starting point. Variations in protocols, secondaries and substrates may require the GM-CSF antibody to be titered up or down for optimal performance.

Immunogen

Recombinant human protein was used as the immunogen for this antibody.

Storage

Store the GM-CSF antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).

References (3)

Ordering:Phone:858.663.9055 | Fax:1.267.821.0800 | Email:info@nsjbio.com

Copyright © NSJ Bioreagents. All rights reserved