


# Connexin 32 Antibody / GJB1 [clone R5.21C] (V3305)

Catalog No.	Formulation	Size
V3305-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3305-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3305SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

V	Citations	(8)	)
		\ - <i>/</i>	,

# **Bulk quote request**

Availability	Discontinued
Species Reactivity	Mouse, Rat
Format	Purified
Clonality	Monoclonal (rat origin)
Isotype	Rat IgG2a, kappa
Clone Name	R5.21C
Purity	Protein G affinity chromatography
UniProt	P28230
Localization	Cell surface, cytoplasmic
Applications	ELISA: 1-5ug/ml for coating (order BSA/sodium azide-free format)
Limitations	This Connexin 32 antibody is available for research use only.



SDS-PAGE Analysis of Purified, BSA-Free Connexin 32 Antibody (clone R5.21C). Confirmation of Integrity and Purity of the Antibody.

## **Description**

This Ab recognizes a protein of 27-32 kDa, identified as Connexin 32. The connexin family of proteins forms hexameric complexes called connexons that facilitate movement of low molecular weight proteins between cells via gap junctions. Connexin proteins share a common topology of four transmembrane α-helical domains, two extracellular loops, a cytoplasmic loop and cytoplasmic N- and C-termini. Many of the key functional differences arise from specific amino-acid substitutions in the most highly conserved domains, the transmembrane and extracellular regions. Each of the approximately 20-connexin isoforms produces channels with distinct permeability and electrical and chemical sensitivities; therefore, one connexin usually cannot fully substitute for another.

# **Application Notes**

Optimal dilution of the Connexin 32 antibody should be determined by the researcher.

### Immunogen

Mouse liver DOC-JR-plasma membranes were used as the immunogen for the Connexin 32 antibody.

### **Storage**

Store the Connexin 32 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).

Ordering:Phone:858.663.9055 | Fax:1.267.821.0800 | Email:info@nsjbio.com

Copyright © NSJ Bioreagents. All rights reserved