


MAML3 Antibody [clone MAML3/1303] (V3531)

Catalog No.	Formulation	Size
V3531-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3531-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3531SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V3531IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml

[Bulk quote request](#)


Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	MAML3/1303
Purity	Protein G affinity chromatography
Buffer	1X PBS, pH 7.4
Gene ID	55534
Localization	Cytoplasmic
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
Limitations	This MAML3 antibody is available for research use only.




IHC testing of FFPE human pancreas with MAML3 antibody (clone MAML3/1303). Staining of FFPE tissue requires boiling sections in 10mM Tris with 1mM EDTA, pH9, for 10-20 min followed by cooling at RT for 20 min.




IHC testing of FFPE human cervical carcinoma with MAML3 antibody (clone MAML3/1303). Staining of FFPE tissue requires boiling sections in 10mM Tris with 1mM EDTA, pH9, for 10-20 min followed by cooling at RT for 20 min.



IHC testing of FFPE human colon carcinoma with MAML3 antibody (clone MAML3/1303). Staining of FFPE tissue requires boiling sections in 10mM Tris with 1mM EDTA, pH9, for 10-20 min followed by cooling at RT for 20 min.



IHC testing of FFPE human placenta with MAML3 antibody (clone MAML3/1303). Staining of FFPE tissue requires boiling sections in 10mM Tris with 1mM EDTA, pH9, for 10-20 min followed by cooling at RT for 20 min.



SDS-PAGE Analysis of Purified, BSA-Free MAML3 Antibody (clone MAML3/1303). Confirmation of Integrity and Purity of the Antibody.

Description

Mastermind-like 3 is a nuclear speckle protein that acts as a transcriptional coactivator for Notch receptors. The Notch signaling pathway influences cell fate by regulating the ability of precursor cells to properly respond to developmental signals. MAML3 is a member of the mastermind-like family of proteins that are human homologs of the *Drosophila melanogaster* mastermind protein. Through its N-terminal region, MAML3 interacts with the ankyrin repeats of the Notch 1-4 proteins. This interaction leads to formation of a DNA-binding complex with the Notch proteins and RBP-J δ ; a complex that can then induce HES1 gene expression. While the N-terminal domain of MAML3 is essential for proper Notch binding, the C-terminal domain

is essential for transcriptional activation. Due to its involvement in cell signaling and transcriptional activation, its upregulation is thought to be involved in oncogenesis.

Application Notes

The concentration stated for each application is a general starting point. Variations in protocols, secondaries and substrates may require the MAML3 antibody to be titrated up or down for optimal performance.

Immunogen

A human full length recombinant protein was used as the immunogen for this MAML3 antibody.

Storage

Store the MAML3 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).

References (1)