


Recombinant Villin Antibody / VIL1 [clone rVIL1/1325] (V3566)

Catalog No.	Formulation	Size
V3566-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3566-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3566SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V3566IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml


Recombinant **MOUSE MONOCLONAL**

[Bulk quote request](#)

Species Reactivity	Human
Format	Purified
Clonality	Recombinant Mouse Monoclonal
Isotype	Mouse IgG1, kappa
Clone Name	rVIL1/1325
Purity	Protein G affinity chromatography
UniProt	P09327
Localization	Cytoplasmic and cell surface
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
Limitations	This recombinant Villin antibody is available for research use only.




IHC testing of FFPE human small intestine carcinoma with recombinant Villin antibody (clone rVIL1/1325). Required HIER: boil tissue sections in pH9 EDTA buffer, for 10-20 min followed by cooling at RT for 20 min.



IHC testing of FFPE human small intestine carcinoma with recombinant Villin antibody (clone rVIL1/1325). Required HIER: boil tissue sections in pH9 EDTA buffer, for 10-20 min followed by cooling at RT for 20 min.

Human Protein Microarray Specificity Validation



Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using recombinant Villin antibody (clone rVIL1/1325). These results demonstrate the foremost specificity of the rVIL1/1325 mAb. Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.

Description

Recognizes a protein of 95kDa, which is identified as villin. It is a major constituent in the microvilli, which compose the brush border of epithelial cells forming absorptive surfaces of the intestinal and renal proximal tubular epithelia. Anti-Villin labels the brush border area in the gastrointestinal mucosal epithelium and urogenital tract. Among neoplasms, villin is predominantly expressed in tumors of colorectal origin. Antibody to villin is useful in identifying malignant cells from primary and metastatic colorectal carcinomas. This antibody also labels Merkel cells of the skin.

Application Notes

The stated application concentrations are suggested starting points. Titration of the recombinant Villin antibody may be required due to differences in protocols and secondary/substrate sensitivity.

1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

Amino acids 179-311 from the human protein were used as the immunogen for this recombinant Villin antibody.

Storage

Store the recombinant Villin antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).