PU.1 Antibody / SPI1 [clone PU1/2146] (V3875)

Catalog No.	Formulation	Size
V3875-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3875-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3875SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2b, kappa
Clone Name	PU1/2146
Purity	Protein G affinity chromatography
UniProt	P17947
Localization	Nuclear
Applications	Flow cytometry : 1-2ug/10^6 cells Immunofluorescence : 1-2ug/ml Western blot : 1-2ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
Limitations	This PU.1 antibody is available for research use only.



IHC staining of FFPE human spleen with PU.1 antibody (clone PU1/2146). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.



IHC staining of FFPE human lymph node with PU.1 antibody (clone PU1/2146). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.



IHC staining of FFPE Hodgkin's lymphoma with PU.1 antibody (clone PU1/2146). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.



IHC staining of FFPE human colon carcinoma with PU.1 antibody (clone PU1/2146). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.



Western blot testing of human THP-1 cell lysate with PU.1 antibody (clone PU1/2146). Predicted molecular weight ~31 kDa but routinely observed at ~40 kDa.



Immunofluorescent staining of PFA-fixed human Ramos cells with PU.1 antibody (green, clone PU1/2146) and Phalloidin (red).



Human Protein Microarray Specificity Validation 100 S score Strength of Signal (Z score) 75 AQP7 3.11 0.39 50 C17orf50 2.72 1.27 25 0 10 30 20 Sig al Ran

Flow cytometry testing of PFA-fixed human Ramos cells with PU.1 antibody (clone PU1/2146); Red=isotype control, Blue= PU.1 antibody.

Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using PU.1 antibody (clone PU1/2146). These results demonstrate the foremost specificity of the PU1/2146 mAb.

Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.



SDS-PAGE analysis of purified, BSA-free PU.1 antibody (clone PU1/2146) as confirmation of integrity and purity.

Description

PU.1 is a member of the ETS family of transcription factors and is important for normal B-cell development. It is expressed in the myeloid lineage and in immature as well as mature B-lymphocytes, with the exception of plasma cells. PU.1 is expressed in germinal center B-cells and mantle B-cells. Various lymphomas are also positive for this marker. It is essential during early B-cell differentiation. The absence of PU.1 results in total block of B-cell development at the pre-pro stage. PU.1 plays a key role in normal myeloid differentiation, and regulates the expression of immunoglobulin and other genes that are important for B-cell development.

Application Notes

Optimal dilution of the PU.1 antibody should be determined by the researcher.

Immunogen

A portion of amino acids 16-170 from the human protein was used as the immunogen for the PU.1 antibody.

Storage

Store the PU.1 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).

Ordering:Phone:858.663.9055 | Fax:1.267.821.0800 | Email:info@nsjbio.com

Copyright © NSJ Bioreagents. All rights reserved