

MECA-79 Antibody / PNAd / Peripheral Node Addressin [clone MECA-79] (V8344)

Catalog No.	Formulation	Size
V8344-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V8344-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V8344SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human, Mouse, Rat
Format	Purified
Clonality	Monoclonal (rat origin)
Isotype	Rat IgM, kappa
Clone Name	MECA-79
Purity	Protein G affinity chromatography
Applications	Flow cytometry : 1-2ug/10 ⁶ cells Immunofluorescence : 1-2ug/ml Western blot : 1-2ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This MECA-79 antibody is available for research use only.

Description

MECA-79 recognizes a carbohydrate epitope shared with a group of sulfation decorated sialomucins, including sulfated ligands for CD62L (CD34, GlyCAM-1, Sgp200, and a subset of MAdCAM-1). This set antigens has been referred to as peripheral node addressin (PNAd) with the molecular mass 50-250 kD. It has been identified that GlcNAc-6-SO₄ sulfation contributes to MECA-79 binding and the core 1 beta1,3-N-acetylglucosaminyltransferase is required for the generation of the MECA-79 epitope. MECA-79 is expressed on high endothelial venules (HEV) of lymphoid tissues, chronic inflamed tissues and rheumatoid synovia. The interaction of PNAd with CD62L receptor is involved in tethering and rolling of lymphocytes along HEV in lymphoid tissues. MECA-79 antibody reacts with Mouse, human and many other species PNAd and blocks L-selectin-dependent lymphocyte adhesion in vitro and in vivo.

Application Notes

Optimal dilution of the MECA-79 antibody should be determined by the researcher.

Immunogen

Mouse lymph node stromal cells were used as the immunogen for this MECA-79 antibody.

Storage

Store the MECA-79 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).