

CATB (Cleaved-Asn126) rabbit pAb

Cat No.: ES19958

For research use only

Overview

Product Name CATB (Cleaved-Asn126) rabbit pAb

Host species Rabbit
Applications WB; ELISA

Species Cross-Reactivity Human;Rat;Mouse;

Recommended dilutions WB 1:1000-2000 ELISA 1:5000-20000

Immunogen Synthesized peptide derived from human CATB

(Cleaved-Asn126)

Specificity This antibody detects endogenous levels of Human

CATB (Cleaved-Asn126, protein was cleaved amino

acid sequence between 126-127)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20° C. Avoid repeated freeze-thaw cycles.

Protein Name CATB (Cleaved-Asn126)

Gene Name CTSB CPSB

Cellular localization Lysosome . Melanosome . Secreted, extracellular

space. Apical cell membrane; Peripheral membrane

protein; Extracellular side. Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:17081065). Localizes to the lumen of thyroid follicles and to the apical

membrane of thyroid epithelial cells (By similarity). .

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 15 37kD
Human Gene ID 1508
Human Swiss-Prot Number P07858

Alternative Names Cathepsin B (EC 3.4.22.1;APP

secretase; APPS; Cathepsin B1) [Cleaved into: Cathepsin B light chain; Cathepsin B heavy chain]



+86-27-59760950 ELKbio@ELKbiotech.com www.e



Background

catalytic activity:Hydrolysis of proteins with broad specificity for peptide bonds. Preferentially cleaves -Arg-Arg-|-Xaa bonds in small molecule substrates (thus differing from cathepsin L). In addition to being an endopeptidase, shows peptidyl-dipeptidase activity, liberating C-terminal dipeptides.,function:Thiol protease which is believed to participate in intracellular degradation and turnover of proteins. Has also been implicated in tumor invasion and metastasis.,similarity:Belongs to the peptidase C1 family.,subcellular location:Identified by mass spectrometry in melanosome fractions from stage I to stage IV.,subunit:Dimer of a heavy chain and a light chain cross-linked by a disulfide bond.,

