

OR3A3 rabbit pAb

Cat No.:ES14377

For research use only

Overview

OR3A3 rabbit pAb
Rabbit
WB
Human;Rat;Mouse;
WB 1: 500-2000
Synthesized peptide derived from human OR3A3 AA range: 256-306
This antibody detects endogenous levels of OR3A3 at Human
Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Store at -20°C. Avoid repeated freeze-thaw cycles.
OR3A3
OR3A3 OR3A6 OR3A7 OR3A8P
Cell membrane; Multi-pass membrane protein.
The antibody was affinity-purified from rabbit
antiserum by affinity-chromatography using
epitope-specific immunogen.
Polyclonal
1 mg/ml
8392
P47888
olfactory receptor family 3 subfamily A member 3(OR3A3) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],

Western blot analysis of lysates from 3T3 cells, primary antibody was diluted at 1:1000, 4° over night





+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C