

Mcl-1 (phospho-Ser64) rabbit pAb

Cat No.: ES15005

For research use only

Overview

Product Name Mcl-1 (phospho-Ser64) rabbit pAb

Host species Rabbit

Applications WB; ELISA;IHC **Species Cross-Reactivity** Human;Rat;Mouse;

Recommended dilutions WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000

Immunogen Synthesized phosho peptide around human Mcl-1

(Ser64)

Specificity This antibody detects endogenous levels of

Human Mcl-1 (phospho-Ser64)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Mcl-1 (Ser64)
Gene Name MCL1 BCL2L3

Cellular localization Membrane; Single-pass membrane protein.

Cytoplasm. Mitochondrion. Nucleus, nucleoplasm.

 $\label{lem:cytoplasmic} \textbf{Cytoplasmic, associated with mitochondria.}$

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal Concentration 1 mg/ml

Observed band About 40kd in human,39kd in mouse and rat

Human Gene ID 4170 Human Swiss-Prot Number Q07820

Alternative Names Induced myeloid leukemia cell differentiation

protein Mcl-1 (Bcl-2-like protein 3) (Bcl2-L-3) (Bcl-2-related protein EAT/mcl1) (mcl1/EAT)

Background This gene encodes an anti-apoptotic protein, which

is a member of the Bcl-2 family. Alternative splicing results in multiple transcript variants. The longest gene product (isoform 1) enhances cell survival by inhibiting apoptosis while the alternatively spliced



+86-27-59760950 ELKbio@ELKbiotech.com


www.elkbiotech.com




shorter gene products (isoform 2 and isoform 3) promote apoptosis and are death-inducing. [provided by RefSeq, Oct 2010],

Western blot analysis of lysates from HCT116 cells, primary antibody was diluted at 1:1000, 4° over night

Immunohistochemical analysis of paraffin-embedded



human Gastric adenocarcinoma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).



+86-27-59760950

