



SPAT2 rabbit pAb


Cat No.:ES12998

For research use only


Overview

| | |
|---------------------------------|---|
| Product Name | SPAT2 rabbit pAb |
| Host species | Rabbit |
| Applications | WB;ELISA;IHC |
| Species Cross-Reactivity | Human;Rat;Mouse; |
| Recommended dilutions | WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000 |
| Immunogen | Synthesized peptide derived from human SPAT2 AA range: 119-169 |
| Specificity | This antibody detects endogenous levels of SPAT2 at Human |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Storage | Store at -20°C. Avoid repeated freeze-thaw cycles. |
| Protein Name | SPAT2 |
| Gene Name | SPATA2 KIAA0757 PD1 |
| Cellular localization | Cytoplasm . Nucleus . Detected in the tubular compartment of the testis and, in the cytoplasm of the Sertoli cells. . |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Clonality | Polyclonal |
| Concentration | 1 mg/ml |
| Observed band | |
| Human Gene ID | 9825 |
| Human Swiss-Prot Number | Q9UM82 |
| Alternative Names | |
| Background | |





Western blot analysis of lysates from MCF-7 cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

