

## Histone H2A (Acetyl Lys9) rabbit pAb

## Cat No.:ES20535

For research use only

## Overview

Product Name	Histone H2A (Acetyl Lys9) rabbit pAb
Host species	Rabbit
Application s	WB
Species Cross-Reac tivity	Human;Mouse;Rat
Recommen ded dilutions	WB: 1:1000-2000
Immunoge n	Synthetic Peptide of Histone H2A (Acetyl Lys9)
Specificity Formulatio n	The antibody detects endogenous Histone H2A (Acetyl Lys9) protein. Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage Protein Name	Store at -20°C. Avoid repeated freeze-thaw cycles. Histone H2A type 1/Histone H2A type 2/Histone H2A type 3
Gene Name	HIST1H2AG/HIST1H2AI/HIST1H2AK/HIST1H2AL/HIST1H2AM/HIST2H2A A3/HIST2H2AA4/HIST3H2A
Cellular localization	Nucleus. Chromosome.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Clonality Concentrat ion	Polyclonal
Observed band	14kD
Human Gene ID	8329/8330/8332/8336/8969/723790/8337/92815



+86-27-59760950

ELKbio@ELKbiotech.com


www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



Human Swiss-Prot Number	POCOS8/Q6FI13/Q7L7L0
Alternative	HIST1H2AG; H2AFP; HIST1H2AI; H2AFC; HIST1H2AK; H2AFD;
Names	HIST1H2AL; H2AFI; HIST1H2AM; H2AFN; Histone H2A type 1; H2A.1; Histone H2A/p; HIST2H2AA3; H2AFO; HIST2H2AA; HIST2H2AA4; Histone H2A type 2-A; Histone H2A.2; Histone H2A/o; HIST3H2A; Histone H2A type 3;
Backgroun	Histones are basic nuclear proteins that are responsible for the
d	nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H2A family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the small histone gene cluster on chromosome 6p22-p21.3. [provided by RefSeq, Aug 2015],

Western blot analysis of extracts from Hela cells, untreated (-) or treated, 1:5000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000





+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C