

## NLTP rabbit pAb

Cat No.: ES14479

For research use only

## Overview

Product Name NLTP rabbit pAb

Host species Rabbit
Applications WB

Species Cross-Reactivity Human; Mouse; Rat Recommended dilutions WB 1: 500-2000

Immunogen Synthesized peptide derived from human NLTP AA

range: 338-388

**Specificity** This antibody detects endogenous levels of NLTP at

Human/Mouse/Rat

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

**Storage** Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name NLTP Gene Name SCP2

**Cellular localization** [Isoform SCP2]: Peroxisome . Cytoplasm .

Mitochondrion . Endoplasmic reticulum .

Mitochondrion .; [Isoform SCPx]: Peroxisome .

**Purification** The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal Concentration 1 mg/ml

**Observed band** 

Human Gene ID 6342 Human Swiss-Prot Number P22307

**Alternative Names** 

**Background** This gene encodes two proteins: sterol carrier

protein X (SCPx) and sterol carrier protein 2 (SCP2),

as a result of transcription initiation from 2

independently regulated promoters. The transcript initiated from the proximal promoter encodes the longer SCPx protein, and the transcript initiated from the distal promoter encodes the shorter SCP2




+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com



protein, with the 2 proteins sharing a common C-terminus. Evidence suggests that the SCPx protein is a peroxisome-associated thiolase that is involved in the oxidation of branched chain fatty acids, while the SCP2 protein is thought to be an intracellular lipid transfer protein. This gene is highly expressed in organs involved in lipid metabolism, and may play a role in Zellweger syndrome, in which cells are deficient in peroxisomes and have impaired bile acid synthesis. Alternative splicing of this gene produces multiple transcript variants, some encoding different isoforms.[provided by RefSeq, Aug 2010],

Western blot analysis of lysates from HuvEc cells, primary antibody was diluted at 1:1000, 4° over night





+86-27-59760950