



SRY (Acetyl Lys136) rabbit pAb

Cat No.:ES20129

For research use only

Overview

Product Name	SRY (Acetyl Lys136) rabbit pAb
Host species	Rabbit
Applications	WB; ELISA
Species Cross-Reactivity	Human;Rat;Mouse;
Recommended dilutions	WB 1:1000-2000 ELISA 1:5000-20000
Immunogen	Synthesized peptide derived from human SRY (Acetyl Lys136)
Specificity	This antibody detects endogenous levels of Human SRY (Acetyl Lys136)
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage	Store at -20°C . Avoid repeated freeze-thaw cycles.
Protein Name	SRY (Acetyl Lys136)
Gene Name	SRY TDF
Cellular localization	Nucleus speckle . Cytoplasm . Nucleus . Acetylation contributes to its nuclear localization and deacetylation by HDAC3 induces a cytoplasmic delocalization (PubMed:15297880). Colocalizes with SOX6 in speckles (PubMed:11818535). Colocalizes with CAML in the nucleus (PubMed:15746192). Colocalizes in the nucleus with ZNF208 isoform KRAB-O and tyrosine hydroxylase (TH) (By similarity). Nuclear import is facilitated by XPO4, a protein that usually acts as a nuclear export signal receptor (PubMed:19349578). .
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	37kD
Human Gene ID	6736
Human Swiss-Prot Number	Q05066





Alternative Names

Background

Sex-determining region Y protein (Testis-determining factor)

disease:Defects in SRY are a cause of gonadal dysgenesis XY female type (GDXY) [MIM:306100]; also known as 'XY females' or Swyer syndrome. Patients are found to have a 46,XY karyotype. They suffer rapid and early degeneration of their gonads, which are present in the adult as 'streak gonads', consisting mainly of fibrous tissue and variable amounts of ovarian stroma. As a result these patients do not develop secondary sexual characteristics at puberty. The external genitalia in these subjects are completely female, and Mullerian structures are normal. In contrast, subjects with 46,XY partial gonadal dysgenesis have ambiguous genitalia, a mix of Mullerian and Wolffian structures, and dysgenetic gonads.

disease:Defects in SRY are a cause of true hermaphroditism [MIM:235600]. A true hermaphrodite must have both mature ovarian and mature testicular tissue with histologic evidence of follicles and tubules, respectively. It is a genetically heterogeneous condition. The genotype of most affected individuals is 46,XX, but many have 46,XY or a mosaic of 46,XX/46,XY. True hermaphroditism can be caused also by chromosomal

translocation.,disease:Defects in SRY are found in Turner syndrome, a disease characterized by gonadal dysgenesis with short stature, "streak gonads", variable abnormalities such as webbing of the neck, cubitus valgus, cardiac defects, low posterior hair line. The phenotype is female. Turner syndrome has its basis in a 45,X chromosomal aberration.

function:Transcriptional regulator which control a genetic switch in male development. It is necessary and sufficient for initiating male sex determination by directing the development of supporting cell precursors (pre-Sertoli cells) as Sertoli rather than granulosa cells (By similarity). In male adult brain involved in the maintenance of motor functions of dopaminergic neurons (By similarity). Involved in different aspects of gene





regulation including promoter activation or repression (By similarity). Facilitates DNA bending. SRY HMG box recognizes DNA by partial intercalation in the minor groove. Also involved in pre-mRNA splicing. Binds to the DNA consensus sequence 5'-[AT]AACAA[AT]-3', miscellaneous: DNA binding and bending properties of the HMG domains of human and mouse SRY differ from each other. Human SRY shows more extensive minor groove contacts with DNA and a lower specificity of sequence recognition than mouse SRY., online information: SRY entry, online information: The tenuous nature of sex - Issue 80 of March 2007, PTM: Phosphorylated on serine residues by PKA. Phosphorylation by PKA enhances its DNA-binding activity and stimulates transcription repression. Acetylation of Lys-136 contributes to its nuclear localization and enhances its interaction with KPNB1. Deacetylated by HDAC3. Poly-ADP-ribosylated by PARP1. ADP-ribosylation reduces its DNA-binding activity., similarity: Belongs to the SRY family., similarity: Contains 1 HMG box DNA-binding domain., subcellular location: Colocalizes with SOX6 in speckles. Colocalizes with CAML in the nucleus. Colocalizes in the nucleus with ZNF208 isoform KRAB-O and tyrosine hydroxylase (TH)., subunit: Interacts with CALM, EP300, HDAC3, KPNB1, ZNF208 isoform KRAB-O, PARP1, SLC9A3R2 and WT1. The interaction with EP300 modulates its DNA-binding activity. The interaction with KPNB1 is sensitive to dissociation by Ran in the GTP-bound form (By similarity). Interaction with PARP1 impaired its DNA-binding activity.,

