

# FAK (phospho Tyr861) rabbit pAb

Cat No.:ES1312

For research use only

## Overview

Product Name	FAK (phospho Tyr861) rabbit pAb
Host species	Rabbit
Applications	WB;ELISA
Species Cross-Reactivity	Human;Mouse;Rat;Monkey
Recommended dilutions	Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized peptide derived from human FAK around the phosphorylation site of Tyr861. AA range:828-877
Specificity	Phospho-FAK (Y861) Polyclonal Antibody detects endogenous levels of FAK protein only when phosphorylated at Y861.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Focal adhesion kinase 1
Gene Name	PTK2
Cellular localization	Cell junction, focal adhesion. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasm, cell cortex. Cytoplasm, cytoskeleton. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Nucleus. Cytoplasm, cytoskeleton, cilium basal body . Constituent of focal adhesions. Detected at microtubules.
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	125kD
Human Gene ID	5747
Human Swiss-Prot Number	Q05397
Alternative Names	PTK2; FAK; FAK1; Focal adhesion kinase 1; FADK 1;






## Background

Focal adhesion kinase-related nonkinase; FRNK;  
Protein phosphatase 1 regulatory subunit 71;  
PPP1R71; Protein-tyrosine kinase 2; p125FAK;  
pp125FAK



protein tyrosine kinase 2(PTK2) Homo sapiens  
This gene encodes a cytoplasmic protein tyrosine kinase which is found concentrated in the focal adhesions that form between cells growing in the presence of extracellular matrix constituents. The encoded protein is a member of the FAK subfamily of protein tyrosine kinases but lacks significant sequence similarity to kinases from other subfamilies. Activation of this gene may be an important early step in cell growth and intracellular signal transduction pathways triggered in response to certain neural peptides or to cell interactions with the extracellular matrix. Several transcript variants encoding different isoforms have been found for this gene, but the full-length nature of only four of them have been determined. [provided by RefSeq, Oct 2015],




Western Blot analysis of various cells using Phospho-FAK (Y861) Polyclonal Antibody diluted at 1:1000



Western blot analysis of 3T3 HELA lysis using Phospho-FAK (Y861) antibody. Antibody was diluted at 1:1000



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using FAK (Phospho-Tyr861) Antibody



Western blot analysis of lysates from HeLa cells, Jurkat cells and COS cells, using FAK (Phospho-Tyr861) Antibody. The lane on the right is blocked with the phospho peptide.

