



# ABCD4 rabbit pAb

Cat No.:ES1567

For research use only

## Overview


<b>Product Name</b>	ABCD4 rabbit pAb
<b>Host species</b>	Rabbit
<b>Applications</b>	WB;IHC;IF;ELISA
<b>Species Cross-Reactivity</b>	Human;Mouse
<b>Recommended dilutions</b>	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications.
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human ABCD4. AA range:111-160
<b>Specificity</b>	ABCD4 Polyclonal Antibody detects endogenous levels of ABCD4 protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Storage</b>	Store at -20°C. Avoid repeated freeze-thaw cycles.
<b>Protein Name</b>	ATP-binding cassette sub-family D member 4
<b>Gene Name</b>	ABCD4
<b>Cellular localization</b>	Endoplasmic reticulum membrane ; Multi-pass membrane protein . Lysosome membrane ; Multi-pass membrane protein . Targeted by LMBRD1 lysosomal chaperone to the lysosomal membrane. .
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Clonality</b>	Polyclonal
<b>Concentration</b>	1 mg/ml
<b>Observed band</b>	70kD
<b>Human Gene ID</b>	5826
<b>Human Swiss-Prot Number</b>	O14678
<b>Alternative Names</b>	ABCD4; PXMP1L; ATP-binding cassette sub-family D member 4; PMP70-related protein; P70R; Peroxisomal membrane protein 1-like; PXMP1-L; Peroxisomal membrane protein 69; PMP69






## Background

The protein encoded by this gene is a member of the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the ALD subfamily, which is involved in peroxisomal import of fatty acids and/or fatty acyl-CoAs in the organelle. All known peroxisomal ABC transporters are half transporters which require a partner half transporter molecule to form a functional homodimeric or heterodimeric transporter. The function of this peroxisomal membrane protein is unknown. However, it is speculated that it may function as a heterodimer for another peroxisomal ABC transporter and, therefore, may modify the adrenoleukodystrophy phenotype. It may also play a role in the process of peroxi



Western Blot analysis of various cells using ABCD4  
Polyclonal Antibody diluted at 1:500




Western Blot analysis of HepG2 cells using ABCD4  
Polyclonal Antibody diluted at 1:500






**ELK Biotechnology**



Western blot analysis of the lysates from HeLa cells using ABCD4 antibody.



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road,Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C