

ATP-citrate synthase rabbit pAb

Cat No.:ES1732

For research use only

Overview

Product Name	ATP-citrate synthase rabbit pAb
Host species	Rabbit
Applications	WB;IF;ELISA
Species Cross-Reactivity	Human; Mouse; Rat; Monkey
Recommended dilutions	Western Blot: 1/500 - 1/2000. Immunofluorescence:
	1/200 - 1/1000. ELISA: 1/10000. Not yet tested in
	other applications.
Immunogen	The antiserum was produced against synthesized
-	peptide derived from human ATP-Citrate Lyase. AA
	range:420-469
Specificity	ATP-citrate synthase Polyclonal Antibody detects
	endogenous levels of ATP-citrate synthase protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	ATP-citrate synthase
Gene Name	ACLY
Cellular localization	Cytoplasm, cytosol.
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	120kD
Human Gene ID	47
Human Swiss-Prot Number	P53396
Alternative Names	ACLY; ATP-citrate synthase; ATP-citrate; pro-S-)-lyase;
	ACL; Citrate cleavage enzyme
Background	ATP citrate lyase(ACLY) Homo sapiens ATP
	citrate lyase is the primary enzyme responsible for
	the synthesis of cytosolic acetyl-CoA in many
	tissues. The enzyme is a tetramer (relative
	molecular weight approximately 440,000) of



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



apparently identical subunits. It catalyzes the formation of acetyl-CoA and oxaloacetate from citrate and CoA with a concomitant hydrolysis of ATP to ADP and phosphate. The product, acetyl-CoA, serves several important biosynthetic pathways, including lipogenesis and cholesterogenesis. In nervous tissue, ATP citrate-lyase may be involved in the biosynthesis of acetylcholine. Multiple transcript variants encoding distinct isoforms have been identified for this gene. [provided by RefSeq, Dec 2014],





ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



Western blot analysis of lysates from COS7 cells, treated with Calyculin 50nM 30', using ATP-Citrate Lyase Antibody. The lane on the right is blocked with the synthesized peptide.



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C