

Aven rabbit pAb

Cat No.:ES1736

For research use only

Overview

Product Name	Aven rabbit pAb
Host species	Rabbit
Applications	WB;IHC;IF;ELISA
Species Cross-Reactivity	Human;Monkey
Recommended dilutions	Western Blot: 1/500 - 1/2000.
	Immunohistochemistry: 1/100 - 1/300.
	Immunofluorescence: 1/200 - 1/1000. ELISA:
	1/20000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized
	peptide derived from human AVEN. AA
	range:301-350
Specificity	Aven Polyclonal Antibody detects endogenous levels
	of Aven protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Cell death regulator Aven
Gene Name	AVEN
Cellular localization	Endomembrane system; Peripheral membrane
	protein. Associated with intracellular membranes.
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	55kD
Human Gene ID	57099
Human Swiss-Prot Number	Q9NQS1
Alternative Names	AVEN; Cell death regulator Aven
Background	function:Protects against apoptosis mediated by
	Apaf-1., subcellular location: Associated with
	intracellular membranes.,subunit:Binds Apaf-1,
	BCL-2 and BAD (Bcl-xl).,tissue specificity:Highly



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



COS7

(kD)

117-85-

48-

34-

26-

19-

AVEN--

LOVO COS

-- 117

-- 85


-- 48

-- 34

-- 26 (kD) expressed in testis, ovary, thymus, prostate, spleen, small intestine, colon, heart, skeletal muscle, liver, kidney and pancreas.,

Western Blot analysis of various cells using Aven Polyclonal Antibody

Western blot analysis of lysates from COS7 and LOVO cells, using AVEN Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human Colon cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C