

Caspase-9 rabbit pAb

Cat No.: ES1858

For research use only

Overview

Immunogen

Product Name Caspase-9 rabbit pAb

Host species Rabbit

WB;IHC;IF;ELISA **Applications Species Cross-Reactivity** Human; Mouse; Rat

Recommended dilutions Western Blot: 1/500 - 1/2000.

Immunohistochemistry: 1/100 - 1/300.

Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications. The antiserum was produced against synthesized

peptide derived from human Caspase 9. AA

range:91-140

Specificity Caspase-9 Polyclonal Antibody detects endogenous

levels of Caspase-9 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Store at -20 $^{\circ}$ C. Avoid repeated freeze-thaw cycles. Storage

Protein Name Caspase9 **Gene Name** CASP9

Cellular localization nucleus, mitochondrion, cytosol, apoptosome, Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal Concentration 1 mg/ml **Observed band** 46kD **Human Gene ID** 842 **Human Swiss-Prot Number** P55211

Alternative Names CASP9; MCH6; Caspase-9; CASP-9; Apoptotic

> protease Mch-6; Apoptotic protease-activating factor 3; APAF-3; ICE-like apoptotic protease 6;

ICE-LAP6

This gene encodes a member of the Background

cysteine-aspartic acid protease (caspase) family.



+86-27-59760950 ELKbio@ELKbiotech.com www.elkbiotech.com



Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. This protein can undergo autoproteolytic processing and activation by the apoptosome, a protein complex of cytochrome c and the apoptotic peptidase activating factor 1; this step is thought to be one of the earliest in the caspase activation cascade. This protein is thought to play a central role in apoptosis and to be a tumor suppressor. Alternative splicing results in multiple transcript variants. [provided by RefSeq, May 2013],



+86-27-59760950