

CNG-1^β rabbit pAb

Cat No.:ES2012

For research use only

Overview

Product Name	CNG-1β rabbit pAb
Host species	Rabbit
Applications	WB;ELISA
Species Cross-Reactivity	Human;Mouse;Rat
Recommended dilutions	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not
	yet tested in other applications.
Immunogen	The antiserum was produced against synthesized
	peptide derived from human CNGB1. AA
	range:571-620
Specificity	CNG-1 ^β Polyclonal Antibody detects endogenous
	levels of CNG-1β protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Cyclic nucleotide-gated cation channel beta-1
Gene Name	CNGB1
Cellular localization	Membrane; Multi-pass membrane protein.
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	102kD
Human Gene ID	1258
Human Swiss-Prot Number	Q14028
Alternative Names	CNGB1; CNCG2; CNCG3L; CNCG4; RCNC2; Cyclic
	nucleotide-gated cation channel beta-1; Cyclic
	nucleotide-gated cation channel 4; CNG channel 4;
	CNG-4; CNG4; Cyclic nucleotide-gated cation
	channel gamma; Cyclic nucleotide-gated cation
	channel mo
Background	In humans, the rod photoreceptor cGMP-gated
	cation channel helps regulate ion flow into the rod



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



KDa 3T3

-CNG-1ß

118

90

50

39

27

138--100--

70---55---40---35---25---

15----

CNGB1--

3T3

CNG-1B

A549 3T3 A549

-- 170

-- 130

-- 95

-- 72

(kD)

photoreceptor outer segment in response to light-induced alteration of the levels of intracellular cGMP. This channel consists of two subunits, alpha and beta, with the protein encoded by this gene representing the beta subunit. Defects in this gene are a cause of cause of retinitis pigmentosa type 45. Three transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2013],

Western Blot analysis of various cells using CNG-1 β Polyclonal Antibody

Western Blot analysis of 3T3 cells using CNG-1 β Polyclonal Antibody

Western blot analysis of lysates from NIH/3T3 and A549 cells, using CNGB1 Antibody. The lane on the right is blocked with the synthesized peptide.



+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C





+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C