

## HSP40 rabbit pAb

Cat No.: ES2583

For research use only

## Overview

Product Name HSP40 rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA Species Cross-Reactivity Human;Mouse;Rat

**Recommended dilutions** Western Blot: 1/500 - 1/2000.

Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human HSP40. AA

range:271-320

Specificity HSP40 Polyclonal Antibody detects endogenous

levels of HSP40 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

**Storage** Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name DnaJ homolog subfamily B member 1

Gene Name DNAJB1

Cellular localization Cytoplasm . Nucleus . Nucleus, nucleolus .

Translocates rapidly from the cytoplasm to the nucleus, and especially to the nucleoli, upon heat

shock.

**Purification** The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 45kD
Human Gene ID 3337
Human Swiss-Prot Number P25685

Alternative Names DNAJB1; DNAJ1; HDJ1; HSPF1; DnaJ homolog

subfamily B member 1; DnaJ protein homolog 1; Heat shock 40 kDa protein 1; HSP40; Heat shock




+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com




**Background** 

protein 40; Human DnaJ protein 1; hDj-1
This gene encodes a member of the DnaJ or Hsp40
(heat shock protein 40 kD) family of proteins. DNAJ
family members are characterized by a highly
conserved amino acid stretch called the 'J-domain'
and function as one of the two major classes of
molecular chaperones involved in a wide range of
cellular events, such as protein folding and
oligomeric protein complex assembly. The encoded
protein is a molecular chaperone that stimulates the
ATPase activity of Hsp70 heat-shock proteins in
order to promote protein folding and prevent
misfolded protein aggregation. Alternative splicing
results in multiple transcript variants. [provided by
RefSeq, Sep 2015],




Immunofluorescence analysis of Hela cell. 1,HSP40 Polyclonal Antibody(red) was diluted at 1:200(4° overnight). Caspase-8 Monoclonal Antibody(2G12)(green) was diluted at 1:200(4° overnight). 2, Goat Anti Rabbit Alexa Fluor 594 Catalog:RS3611 was diluted at




Western Blot analysis of various cells using HSP40 Polyclonal Antibody










Immunohistochemistry analysis of paraffin-embedded human lung carcinoma tissue, using HSP40 Antibody. The picture on the right is blocked with the synthesized peptide.



+86-27-59760950

