

MEF-2 rabbit pAb

Cat No.:ES2766

For research use only

Overview

Product Name MEF-2 rabbit pAb

Host species Rabbit

Applications WB;IHC;IF;ELISA Species Cross-Reactivity Human;Mouse;Rat

Recommended dilutions Western Blot: 1/500 - 1/2000.

Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human MEF2A. AA

range:374-423

Specificity MEF-2 Polyclonal Antibody detects endogenous

levels of MEF-2 protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Myocyte-specific enhancer factor 2A

Gene Name MEF2A **Cellular localization** Nucleus .

Purification The antibody was affinity-purified from rabbit

antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 55kD
Human Gene ID 4205
Human Swiss-Prot Number Q02078

Alternative Names MEF2A; MEF2; Myocyte-specific enhancer factor 2A;

Serum response factor-like protein 1

Background The protein encoded by this gene is a DNA-binding


transcription factor that activates many muscle-specific, growth factor-induced, and

stress-induced genes. The encoded protein can act







as a homodimer or as a heterodimer and is involved in several cellular processes, including muscle development, neuronal differentiation, cell growth control, and apoptosis. Defects in this gene could be a cause of autosomal dominant coronary artery disease 1 with myocardial infarction (ADCAD1). Several transcript variants encoding different isoforms have been found for this gene.[provided by RefSeq, Jan 2010],



Western Blot analysis of various cells using MEF-2 Polyclonal Antibody diluted at 1:1000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).



Immunofluorescence analysis of HeLa cells, using MEF2A Antibody. The picture on the right is blocked with the synthesized peptide.




Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using MEF2A Antibody. The picture on the right is blocked with the synthesized peptide.



+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com





Western blot analysis of lysates from HeLa cells, treated with PMA 125ng/ml 30', using MEF2A Antibody. The lane on the right is blocked with the synthesized peptide.

