

Olfactory receptor 11G2 rabbit pAb

Cat No.:ES3017

For research use only

Overview

Product Name	Olfactory receptor 11G2 rabbit pAb
Host species	Rabbit
Applications	WB;IF;ELISA
Species Cross-Reactivity	Human;Rat;Mouse;
Recommended dilutions	Western Blot: 1/500 - 1/2000. Immunofluorescence:
	1/200 - 1/1000. ELISA: 1/40000. Not yet tested in
	other applications.
Immunogen	The antiserum was produced against synthesized
	peptide derived from human OR11G2. AA
	range:241-290
Specificity	Olfactory receptor 11G2 Polyclonal Antibody detects
	endogenous levels of Olfactory receptor 11G2
	protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Olfactory receptor 11G2
Gene Name	OR11G2
Cellular localization	Cell membrane; Multi-pass membrane protein.
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	35kD
Human Gene ID	390439
Human Swiss-Prot Number	Q8NGC1
Alternative Names	OR11G2; Olfactory receptor 11G2; Olfactory
	receptor OR14-34
Background	Olfactory receptors interact with odorant molecules
	in the nose, to initiate a neuronal response that
	triggers the perception of a smell. The olfactory
	receptor proteins are members of a large family of



+86-27-59760950

ELKbio@ELKbiotech.com


www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],

Western Blot analysis of various cells using Olfactory receptor 11G2 Polyclonal Antibody diluted at 1:1000





HuvEc

(kD)

117-85-

48-

34-

26-

19-




+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



Western blot analysis of lysates from HUVEC, Jurkat, and MCF-7 cells, using OR11G2 Antibody. The lane on the right is blocked with the synthesized peptide.

Western blot analysis of the lysates from HeLa cells using OR11G2 antibody.



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C