

## Olfactory receptor 51D1 rabbit pAb

Cat No.: ES3049

For research use only

## Overview

Product Name Olfactory receptor 51D1 rabbit pAb

Host species Rabbit
Applications WB;IF;ELISA

Species Cross-Reactivity Human; Rat; Mouse;

**Recommended dilutions** Western Blot: 1/500 - 1/2000. Immunofluorescence:

1/200 - 1/1000. ELISA: 1/10000. Not yet tested in

other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human OR51D1. AA

range:33-82

**Specificity** Olfactory receptor 51D1 Polyclonal Antibody detects

endogenous levels of Olfactory receptor 51D1

protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

**Storage** Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Olfactory receptor 51D1

Gene Name OR51D1

Cellular localizationCell membrane; Multi-pass membrane protein.PurificationThe antibody was affinity-purified from rabbit<br/>antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 34kD
Human Gene ID 390038
Human Swiss-Prot Number Q8NGF3

Alternative Names OR51D1; Olfactory receptor 51D1; Olfactory

receptor OR11-14

**Background** Olfactory receptors interact with odorant molecules


in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of




+86-27-59760950 ELKbio@ELKbiotech.com www



G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],




Western Blot analysis of various cells using Olfactory receptor 51D1 Polyclonal Antibody



Western blot analysis of lysates from Jurkat cells, using OR51D1 Antibody. The lane on the right is blocked with the synthesized peptide.







Western blot analysis of the lysates from COLO205 cells using OR51D1 antibody.

