

Olfactory receptor 6C3 rabbit pAb

Cat No.: ES3067

For research use only

Overview

Product Name Olfactory receptor 6C3 rabbit pAb

Host species Rabbit
Applications WB;IF;ELISA

Species Cross-Reactivity Human; Rat; Mouse;

Recommended dilutions Western Blot: 1/500 - 1/2000. Immunofluorescence:

1/200 - 1/1000. ELISA: 1/10000. Not yet tested in

other applications.

Immunogen The antiserum was produced against synthesized

peptide derived from human OR6C3. AA

range:228-277

Specificity Olfactory receptor 6C3 Polyclonal Antibody detects

endogenous levels of Olfactory receptor 6C3

protein.

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and

0.02% sodium azide.

Storage Store at -20°C. Avoid repeated freeze-thaw cycles.

Protein Name Olfactory receptor 6C3

Gene Name OR6C3

Cellular localizationCell membrane; Multi-pass membrane protein.PurificationThe antibody was affinity-purified from rabbit
antiserum by affinity-chromatography using

epitope-specific immunogen.

Clonality Polyclonal
Concentration 1 mg/ml
Observed band 35kD
Human Gene ID 254786
Human Swiss-Prot Number O9NZP0

Alternative Names OR6C3; Olfactory receptor 6C3; HSA8

Background olfactory receptor family 6 subfamily C member

3(OR6C3) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor




+86-27-59760950 ELKbio@ELKbiotech.com

www.elkbiotech.com




proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],

Western Blot analysis of various cells using Olfactory receptor 6C3 Polyclonal Antibody




Western blot analysis of lysates from HeLa and HepG2 cells, using OR6C3 Antibody. The lane on the right is blocked with the synthesized peptide.



+86-27-59760950







Western blot analysis of the lysates from K562 cells using OR6C3 antibody.

