

Olfactory receptor 6K2 rabbit pAb

Cat No.:ES3069

For research use only

Overview

Product Name	Olfactory receptor 6K2 rabbit pAb
Host species	Rabbit
Applications	WB;ELISA
Species Cross-Reactivity	Human;Rat;Mouse;
Recommended dilutions	Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized
	peptide derived from human OR6K2. AA range:271-320
Specificity	Olfactory receptor 6K2 Polyclonal Antibody detects endogenous levels of Olfactory receptor 6K2
	protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Olfactory receptor 6K2
Gene Name	OR6K2
Cellular localization	Cell membrane; Multi-pass membrane protein.
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	37kD
Human Gene ID	81448
Human Swiss-Prot Number	Q8NGY2
Alternative Names	OR6K2; Olfactory receptor 6K2; Olfactory receptor
	OR1-17
Background	olfactory receptor family 6 subfamily K member
	2(OR6K2) Homo sapiens Olfactory receptors
	interact with odorant molecules in the nose, to
	initiate a neuronal response that triggers the
	perception of a smell. The olfactory receptor



+86-27-59760950

ELKbio@ELKbiotech.com


www.elkbiotech.com


23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],

Western Blot analysis of various cells using Olfactory receptor 6K2 Polyclonal Antibody





HeLa

(kD)




+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



Western blot analysis of the lysates from HepG2 cells using OR6K2 antibody.



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C