

PDXDC1 rabbit pAb

Cat No.:ES3172

For research use only

Overview

Product Name	PDXDC1 rabbit pAb
Host species	Rabbit
Applications	WB;IHC;IF;ELISA
Species Cross-Reactivity	Human;Mouse;Rat
Recommended dilutions	Western Blot: 1/500 - 1/2000.
	Immunohistochemistry: 1/100 - 1/300. ELISA:
	1/10000. Not yet tested in other applications.
Immunogen	The antiserum was produced against synthesized
	peptide derived from human PDXDC1. AA
	range:677-726
Specificity	PDXDC1 Polyclonal Antibody detects endogenous
	levels of PDXDC1 protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and
	0.02% sodium azide.
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.
Protein Name	Pyridoxal-dependent decarboxylase
	domain-containing protein 1
Gene Name	PDXDC1
Cellular localization	Golgi apparatus, cell-cell adherens
	junction, intracellular membrane-bounded organelle,
Purification	The antibody was affinity-purified from rabbit
	antiserum by affinity-chromatography using
	epitope-specific immunogen.
Clonality	Polyclonal
Concentration	1 mg/ml
Observed band	87kD
Human Gene ID	23042
Human Swiss-Prot Number	Q6P996
Alternative Names	PDXDC1; KIAA0251; Pyridoxal-dependent
	decarboxylase domain-containing protein 1
Background	cofactor:Pyridoxal phosphate.,PTM:Phosphorylated
	upon DNA damage, probably by ATM or
	ATR., similarity: Belongs to the group II decarboxylase



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



HepG2

(kD)

170-

130-

95-

72-

55-

PDXDC1 --

(kD)

117-

85-

48-

34-

26-

19-

HepG2 HUVEC HepG2

-- 117

-- 85

-- 48

-- 34

-- 26 -- 19 (kD)

PDXDC1

family.,

Western Blot analysis of various cells using PDXDC1 Polyclonal Antibody


Western blot analysis of lysates from HepG2 and HUVEC cells, using PDXDC1 Antibody. The lane on the right is blocked with the synthesized peptide.

Western blot analysis of the lysates from HeLa cells using PDXDC1 antibody.



23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C





Immunohistochemical analysis of paraffin-embedded human Breast cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C