

## **RUNX3** rabbit pAb

## Cat No.:ES3400

For research use only

## Overview

Product Name	RUNX3 rabbit pAb	
Host species	Rabbit	
Applications	WB;ELISA	
Species Cross-Reactivity	Human;Mouse	
<b>Recommended dilutions</b>	Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not	
	yet tested in other applications.	
Immunogen	The antiserum was produced against synthesized	
	peptide derived from human RUNX3. AA	
	range:133-182	
Specificity	RUNX3 Polyclonal Antibody detects endogenous	
	levels of RUNX3 protein.	
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and	
	0.02% sodium azide.	
Storage	Store at -20°C. Avoid repeated freeze-thaw cycles.	
Protein Name	Runt-related transcription factor 3	
Gene Name	RUNX3	
Cellular localization	Nucleus . Cytoplasm . The tyrosine phosphorylated	
	form localizes to the cytoplasm. Translocates from	
	the cytoplasm to the nucleus following TGF-beta	
	stimulation	
Purification	The antibody was affinity-purified from rabbit	
	antiserum by affinity-chromatography using	
	epitope-specific immunogen.	
Clonality	Polyclonal	
Concentration	1 mg/ml	
Observed band	44kD	
Human Gene ID	864	
Human Swiss-Prot Number	Q13761	
Alternative Names	RUNX3; AML2; CBFA3; PEBP2A3; Runt-related	
	transcription factor 3; Acute myeloid leukemia 2	
	protein; Core-binding factor subunit alpha-3;	
	CBF-alpha-3; Oncogene AML-2; Polyomavirus	
	enhancer-binding protein 2 alpha C subunit;	



+86-27-59760950

ELKbio@ELKbiotech.com

www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C



Background

## PEA2-alpha C; PEB

This gene encodes a member of the runt domain-containing family of transcription factors. A heterodimer of this protein and a beta subunit forms a complex that binds to the core DNA sequence 5'-PYGPYGGT-3' found in a number of enhancers and promoters, and can either activate or suppress transcription. It also interacts with other transcription factors. It functions as a tumor suppressor, and the gene is frequently deleted or transcriptionally silenced in cancer. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Mar 2016],

Western Blot analysis of various cells using RUNX3 Polyclonal Antibody diluted at 1:500 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).







www.elkbiotech.com

23-2, No.388 Gaoxin 2nd Road, Wuhan East Lake Hi-tech Development Zone, Hubei , P.R.C